

# Catch-Up Guidance for Children 4 Months through 17 Years of Age

## Inactivated Polio Vaccine (IPV)<sup>1</sup>

The table below provides guidance for children whose vaccinations have been delayed. Start with the child's age and information on previous doses (previous doses must be documented and must meet minimum age requirements and minimum intervals between doses). Use this table in conjunction with table 2 of the Recommended Child and Adolescent Immunization Schedule for Ages 18 Years or Younger, found at [downloads.aap.org/AAP/PDF/AAP-Immunization-Schedule.pdf](https://downloads.aap.org/AAP/PDF/AAP-Immunization-Schedule.pdf)

IF current age is	AND # of previous doses <sup>2</sup> is	AND		THEN	Next dose due <sup>3</sup>
4 through 18 months	Unknown or 0	→		Give Dose 1 today	Give Dose 2 at least 4 weeks after Dose 1
	1	It has been at least 4 weeks since Dose 1		Give Dose 2 today	Give Dose 3 at least 4 weeks after Dose 2 <b>and</b> at 6 months of age or older
		It has <b>not</b> been at least 4 weeks since Dose 1		No dose today	Give Dose 2 at least 4 weeks after Dose 1
	2	It has been at least 4 weeks since Dose 2	Child is 6 months of age or older	Give Dose 3 today	Give Dose 4 ( <b>Final Dose</b> ) at 4 through 6 years of age <sup>4</sup>
			Child is younger than 6 months of age	No dose today	Give Dose 3 at 6 months of age
		It has <b>not</b> been at least 4 weeks since Dose 2	→	No dose today	Give Dose 3 at least 4 weeks after Dose 2 <b>and</b> at 6 months of age or older
	19 months through 3 years	Unknown or 0	→		Give Dose 1 today
1		It has been at least 4 weeks since Dose 1		Give Dose 2 today	Give Dose 3 at least 4 weeks after Dose 2
		It has <b>not</b> been at least 4 weeks since Dose 1		No dose today	Give Dose 2 at least 4 weeks after Dose 1
2		It has been at least 4 weeks since Dose 2		Give Dose 3 today	Give Dose 4 ( <b>Final Dose</b> ) at least 6 months after Dose 3 and at 4 through 6 years of age <sup>4</sup>
		It has <b>not</b> been 4 weeks since Dose 2		No dose today	Give Dose 3 at least 4 weeks after Dose 2

<sup>1</sup>IPV products include IPOL, Pediarix, Pentacel, Vaxelis, Kinrix, and Quadracel. Use the correct product based on the approved age indications.

<sup>2</sup>Series containing oral polio vaccine (OPV) administered before April 1, 2016, either mixed OPV-IPV or OPV only: Total number of doses needed to complete the series is the same as that recommended for the U.S. IPV schedule. [cdc.gov/mmwr/volumes/66/wr/mm6601a6.htm](https://www.cdc.gov/mmwr/volumes/66/wr/mm6601a6.htm)

<sup>3</sup>Next dose due is not the final dose in the series unless explicitly stated.

<sup>4</sup>Vaxelis is not indicated for Dose 4.



# Catch-Up Guidance for Children 4 Months through 17 Years of Age

## Inactivated Polio Vaccine (IPV)<sup>1</sup>

IF current age is	AND # of previous doses <sup>2</sup> is	AND			THEN	Next dose due <sup>3</sup>	
4 through 17 years	Unknown or 0	→			Give Dose 1 today	Give Dose 2 at least 4 weeks after Dose 1	
	1	It has been at least 4 weeks since Dose 1			Give Dose 2 today	Give Dose 3 ( <b>Final Dose</b> ) at least 6 months after Dose 2 <sup>5</sup>	
		It has <b>not</b> been 4 weeks since Dose 1			No dose today	Give Dose 2 at least 4 weeks after Dose 1	
	2	It has been at least 6 months since Dose 2			Give Dose 3 ( <b>Final Dose</b> ) today <sup>5</sup>	<b>No additional doses needed</b>	
		It has <b>not</b> been 6 months since Dose 2			No dose today	Give Dose 3 ( <b>Final Dose</b> ) at least 6 months after Dose 2 <sup>5</sup>	
	3	Dose 3 was given before 4 years of age	It has been at least 6 months since Dose 3	→	Give Dose 4 ( <b>Final dose</b> ) today <sup>4</sup>	<b>No additional doses needed</b>	
			It has <b>not</b> been at least 6 months since Dose 3	→	No dose today	Give Dose 4 ( <b>Final Dose</b> ) at least 6 months after Dose 3 <sup>4</sup>	
		Dose 3 was given at 4 years of age or older	Dose 3 was given at least 6 months from previous dose	→	No dose today	<b>No additional doses needed</b>	
			Dose 3 was <b>not</b> given at least 6 months from previous dose	It has been at least 6 months since Dose 3	→	Give Dose 4 ( <b>Final dose</b> ) today <sup>4</sup>	<b>No additional doses needed</b>
				It has not been at least 6 months since Dose 3	→	No dose today	Give Dose 4 ( <b>Final Dose</b> ) at least 6 months after Dose 3 <sup>4</sup>

<sup>1</sup>IPV products include IPOL, Pediarix, Pentacel, Vaxelis, Kinrix, and Quadracel. Use the correct product based on the approved age indications.

<sup>2</sup>Series containing oral polio vaccine (OPV) administered before April 1, 2016, either mixed OPV-IPV or OPV only: Total number of doses needed to complete the series is the same as that recommended for the U.S. IPV schedule. [www.cdc.gov/mmwr/volumes/66/wr/mm6601a6.htm](http://www.cdc.gov/mmwr/volumes/66/wr/mm6601a6.htm)

<sup>3</sup>Next dose due is not the final dose in the series unless explicitly stated.

<sup>4</sup>Vaxelis is not indicated for Dose 4.

<sup>5</sup>Dose 4 is not necessary if the Dose 3 was given at age 4 years or older and at least 6 months after the previous dose.

