

# Expanded BRFSS, 2003

## Locality Summary Tables

### Table of Contents:

Optional Module 1: Adult Asthma History .....	1
Optional Module 3: Cardiovascular Disease .....	4
Optional Module 4: Cholesterol Awareness .....	7
Optional Module 5: Diabetes .....	9
Optional Module 6: Disability .....	13
Optional Module 7: Quality of Life .....	14
Optional Module 9: Fruits and Vegetables .....	17
Optional Module 10: Health Care Coverage .....	18
Optional Module 11: HIV/AIDS .....	20
Optional Module 12: Hypertension Awareness .....	21
Optional Module 13: Injury Control - Falls .....	22
Optional Module 14: Immunization .....	23
Optional Module 15: Oral Health .....	27
Optional Module 16: Physical Activity .....	29
Optional Module 17: Skin Cancer .....	34
Optional Module 18: Social Context .....	36
Optional Module 19: Tobacco Consumption .....	40
Optional Module 20: Tobacco - Media .....	41
Optional Module 21: Tobacco – other products .....	49
Optional Module 22: Tobacco – Work site ETS .....	50
Optional Module 23: Seat Belts .....	52
Optional Module 24: Weight Control .....	54
Optional Module 25: Cervical Cancer .....	58

**New York State Department of Health**

Expanded BRFSS, 2003 - New York State Department of Health

**Have had an Asthma Episode or Attack in the Past 12 Months, among Those with Asthma**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Allegany, Wyoming	28	53.1	22	46.9	16.6
Dutchess, Putnam	26	43.6	29	56.4	14.2
Bronx	36	55.0	33	45.0	14.0
Kings (Brooklyn)	36	64.6	20	35.4	14.1
New York (Manhattan)	32	64.1	18	35.9	16.1
Queens	20	41.3	23	58.7	17.5
Richmond (Staten Island)	32	61.8	18	38.2	15.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

This page left blank intentionally

This page left blank intentionally

Expanded BRFSS, 2003 - New York State Department of Health

**Adults Eating Fewer High Fat or High Cholesterol Foods to Lower Risk of Developing Heart Disease or Stroke**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Monroe	396	66.9	202	33.1	4.2
Niagara	398	63.0	213	37.0	4.4

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Adults Eating More Fruits and Vegetables to Lower Risk of Developing Heart Disease or Stroke**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Monroe	413	70.0	194	30.0	4.1
Niagara	410	69.0	206	31.0	4.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Adults More Physically Active to Lower Risk of Developing Heart Disease or Stroke**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Monroe	491	81.9	117	18.1	3.4
Niagara	504	79.8	116	20.2	3.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Ever had Blood Cholesterol Checked**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Albany	518	82.5	89	17.5	3.8
Allegany, Wyoming	472	73.3	133	26.7	4.1
Broome	496	76.4	115	23.6	4.1
Cattaraugus, Chautauqua	497	77.8	107	22.2	4.2
Cayuga, Seneca, Wayne	521	81.0	97	19.0	4.0
Chemung, Schuylar, Tioga	490	77.7	118	22.3	3.9
Chenango, Madison	505	75.8	129	24.2	4.2
Clinton, Essex, Franklin	481	71.9	131	28.1	4.7
Columbia, Greene	522	84.0	80	16.0	3.6
Cortland, Tompkins	496	73.7	125	26.3	4.5
Dutchess, Putnam	518	84.6	78	15.4	3.8
Fulton, Montgomery	493	80.2	115	19.8	3.8
Genesee, Orleans	495	78.3	116	21.7	3.9
Hamilton, Herkimer	494	80.4	93	19.6	4.1
Jefferson, Lewis, StLawrence	512	80.1	98	19.9	4.3
Livingston, Ontario	502	78.4	104	21.6	4.0
Niagara	518	81.8	83	18.2	3.9
Oneida, Oswego	515	77.9	118	22.1	3.9
Orange	537	82.5	84	17.5	4.0
Rensselaer	531	83.6	82	16.4	3.5
Rockland	529	85.4	68	14.6	3.7
Saratoga	545	86.0	71	14.0	3.5
Schenectady	516	81.7	92	18.3	3.8
Steuben, Yates	507	77.8	118	22.2	4.1
Suffolk	542	85.4	75	14.6	3.5
Westchester	505	82.3	80	17.7	4.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health  
**Had Blood Cholesterol Checked in the Past Five Years**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Albany	488	78.8	112	21.2	4.0
Allegany, Wyoming	435	69.3	158	30.7	4.3
Broome	456	71.3	150	28.7	4.3
Cattaraugus, Chautauqua	465	73.0	130	27.0	4.4
Cayuga, Seneca, Wayne	482	75.7	127	24.3	4.3
Chemung, Schuylar, Tioga	453	72.7	149	27.3	4.1
Chenango, Madison	465	70.5	162	29.5	4.4
Clinton, Essex, Franklin	443	65.8	164	34.2	4.8
Columbia, Greene	483	79.7	113	20.3	3.8
Cortland, Tompkins	447	67.0	166	33.0	4.7
Dutchess, Putnam	475	79.5	107	20.5	4.1
Fulton, Montgomery	448	74.3	149	25.7	4.1
Genesee, Orleans	456	72.8	148	27.2	4.2
Hamilton, Herkimer	465	76.1	117	23.9	4.3
Jefferson, Lewis, StLawrence	475	74.4	129	25.6	4.7
Livingston, Ontario	480	75.9	121	24.1	4.1
Niagara	487	77.5	106	22.5	4.2
Oneida, Oswego	476	72.8	151	27.2	4.1
Orange	514	80.0	100	20.0	4.1
Rensselaer	504	79.9	106	20.1	3.7
Rockland	498	80.9	91	19.1	4.0
Saratoga	514	81.7	97	18.3	3.8
Schenectady	492	78.4	112	21.6	4.0
Steuben, Yates	481	72.9	142	27.1	4.5
Suffolk	509	81.0	101	19.0	3.8
Westchester	479	78.7	102	21.3	4.4

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

## Expanded BRFSS, 2003 - New York State Department of Health

**Check Blood for Glucose or Sugar at Least Once per Day among Those with Diabetes**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Allegany, Wyoming	29	55.9	19	44.1	15.5
Dutchess, Putnam	25	58.1	20	41.9	15.8
Fulton, Montgomery	28	54.0	25	46.0	15.0
Niagara	30	53.9	27	46.1	14.5
Onondaga	32	70.6	13	29.4	14.5
Warren, Washington	29	56.2	20	43.8	16.0
Bronx	26	35.3	40	64.7	13.6
Kings (Brooklyn)	25	50.1	24	49.9	15.7
New York (Manhattan)	15	46.9	18	53.1	20.0
Queens	22	39.9	26	60.1	15.5
Richmond (Staten Island)	26	69.0	15	31.0	15.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Check Feet for Sores or Irritations at Least Once per Day among Those with Diabetes**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Allegany, Wyoming	31	59.3	18	40.7	15.1
Dutchess, Putnam	30	72.0	14	28.0	13.9
Fulton, Montgomery	35	65.9	16	34.1	15.4
Niagara	37	67.6	18	32.4	14.1
Onondaga	29	66.3	13	33.7	15.8
Warren, Washington	27	54.8	20	45.2	16.2
Bronx	42	65.2	21	34.8	14.2
Kings (Brooklyn)	25	51.1	23	48.9	15.9
New York (Manhattan)	14	37.3	18	62.7	18.8
Queens	31	57.6	18	42.4	16.0
Richmond (Staten Island)	25	61.6	13	38.4	17.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Had Feet Checked by a Medical Professional in Past Twelve Months, among Those with Diabetes**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Allegany, Wyoming	34	68.0	15	32.0	14.4
Dutchess, Putnam	34	80.6	9	19.4	12.4
Fulton, Montgomery	37	77.0	11	23.0	13.0
Niagara	39	73.6	15	26.4	14.0
Onondaga	34	78.1	9	21.9	13.2
Warren, Washington	36	76.9	10	23.1	14.0
Bronx	47	77.6	15	22.4	12.1
Kings (Brooklyn)	34	75.0	11	25.0	14.7
New York (Manhattan)	21	63.0	10	37.0	21.0
Queens	31	56.2	16	43.8	16.6
Richmond (Staten Island)	27	65.6	12	34.4	17.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution<sup>2</sup>Weighted Percent<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Had Dilated Eye Exam in Past 12 Months, among Those with Diabetes**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Allegany, Wyoming	37	75.2	11	24.8	13.7
Dutchess, Putnam	35	80.2	9	19.8	13.3
Fulton, Montgomery	30	61.2	21	38.8	15.2
Niagara	47	86.6	7	13.4	9.9
Onondaga	32	73.9	13	26.1	13.5
Warren, Washington	32	67.0	16	33.0	14.8
Bronx	49	77.5	14	22.5	12.5
Kings (Brooklyn)	32	70.8	14	29.2	15.4
New York (Manhattan)	30	97.2	2	2.8	4.1
Queens	32	73.5	14	26.5	14.5
Richmond (Staten Island)	30	74.5	11	25.5	14.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Adults with a Disability**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Albany	131	19.2	495	80.8	3.5
Broome	160	22.4	468	77.6	3.6
Cattaraugus, Chautauqua	156	23.0	462	77.0	3.7
Cayuga, Seneca, Wayne	139	19.5	498	80.5	3.4
Chemung, Schuyler, Tioga	141	20.1	481	79.9	3.4
Chenango, Madison	153	19.5	493	80.5	3.2
Clinton, Essex, Franklin	148	20.7	472	79.3	3.5
Columbia, Greene	134	19.5	486	80.5	3.4
Cortland, Tompkins	133	17.4	497	82.6	3.2
Erie	122	18.5	487	81.5	3.3
Genesee, Orleans	152	23.8	471	76.2	3.7
Hamilton, Herkimer	152	23.0	445	77.0	3.8
Jefferson, Lewis, StLawrence	140	17.9	485	82.1	3.4
Livingston, Ontario	139	19.7	481	80.3	3.4
Niagara	138	20.0	481	80.0	3.4
Oneida, Oswego	141	17.9	499	82.1	3.1
Onondaga	106	15.2	518	84.8	3.1
Orange	131	17.9	494	82.1	3.2
Rensselaer	137	19.3	480	80.7	3.3
Rockland	110	16.1	495	83.9	3.2
Saratoga	136	19.9	492	80.1	3.5
Schenectady	140	19.0	484	81.0	3.3
Steuben, Yates	152	22.8	485	77.2	3.9
Westchester	103	14.0	497	86.0	2.9
Bronx	118	17.3	494	82.7	3.4
Kings (Brooklyn)	108	16.3	513	83.7	3.2
New York (Manhattan)	104	14.3	500	85.7	3.0
Queens	96	13.9	487	86.1	3.0
Richmond (Staten Island)	96	13.5	518	86.5	2.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Adults with a Disability Requiring Help**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Disability need help %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Disability no help needed %</b>	<b>C.I.</b>	<b>n</b>	<b>No Disability %</b>	<b>C.I.</b>
Onondaga	37	4.9	1.7	69	10.3	2.7	518	84.8	3.1
Bronx	61	9.5	2.6	57	7.8	2.3	494	82.7	3.4
Kings (Brooklyn)	39	5.3	1.8	69	11.0	2.8	513	83.7	3.2
New York (Manhattan)	41	6.3	2.2	63	8.0	2.1	500	85.7	3.0
Queens	38	5.6	1.9	58	8.3	2.4	487	86.1	3.0
Richmond (Staten Island)	43	6.2	2.0	53	7.3	2.1	518	86.5	2.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution<sup>2</sup>Weighted Percent<sup>3</sup>95% Confidence Interval

This page left intentionally blank



This page left intentionally blank

Expanded BRFSS, 2003 - New York State Department of Health  
**Eat Five Fruits or Vegetable Servings per Day**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Albany	195	29.6	436	70.4	3.9
Allegany, Wyoming	155	22.4	465	77.6	3.6
Broome	167	27.6	461	72.4	4.0
Cattaraugus, Chautauqua	162	23.5	459	76.5	3.7
Cayuga, Seneca, Wayne	163	24.0	474	76.0	3.8
Chemung, Schuylar, Tioga	151	23.0	473	77.0	3.6
Chenango, Madison	149	20.6	501	79.4	3.5
Clinton, Essex, Franklin	154	24.4	470	75.6	4.0
Columbia, Greene	192	29.3	432	70.7	4.0
Cortland, Tompkins	192	27.0	441	73.0	3.9
Delaware, Otsego, Schoharie	199	30.5	420	69.5	4.2
Dutchess, Putnam	184	28.6	420	71.4	4.0
Erie	162	24.5	443	75.5	3.7
Fulton, Montgomery	153	24.5	467	75.5	3.8
Genesee, Orleans	151	23.1	473	76.9	3.6
Hamilton, Herkimer	157	27.1	442	72.9	4.3
Jefferson, Lewis, StLawrence	183	30.3	444	69.7	4.6
Livingston, Ontario	180	29.3	441	70.7	4.1
Monroe	194	31.9	416	68.1	4.1
Niagara	186	29.1	427	70.9	4.0
Oneida, Oswego	164	23.8	478	76.2	3.6
Orange	180	25.2	450	74.8	3.9
Rensselaer	157	24.9	465	75.1	3.8
Rockland	170	26.9	441	73.1	4.0
Saratoga	159	23.4	471	76.6	3.8
Schenectady	191	28.5	432	71.5	4.1
Steuben, Yates	148	22.9	491	77.1	3.7
Suffolk	182	28.7	441	71.3	4.2
Warren, Washington	166	24.7	468	75.3	3.8
Westchester	194	31.6	400	68.4	4.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

**Main Reason for Not Having Health Care Coverage**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Changed or lost job %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Could not afford %</b>	<b>C.I.</b>	<b>n</b>	<b>Other %</b>	<b>C.I.</b>
Allegany, Wyoming	12	22.1	12.7	27	44.4	14.5	21	33.5	13.6
Delaware, Otsego, Schoharie	8	7.0	5.7	36	49.8	15.2	25	43.3	15.4
Dutchess, Putnam	6	13.2	11.0	20	36.5	15.4	25	50.3	16.2
Erie	11	20.7	12.7	9	25.6	16.7	17	53.7	18.5
Fulton, Montgomery	15	24.9	12.7	21	35.7	13.7	25	39.5	13.8
Niagara	9	18.6	13.6	13	31.7	17.3	15	49.8	19.4
Onondaga	13	14.5	8.8	20	34.7	14.5	22	50.8	15.7
Suffolk	8	13.9	9.7	23	33.1	13.9	23	53.0	16.0
Sullivan, Ulster	17	22.8	11.1	41	39.6	12.4	24	37.6	13.3
Westchester	15	18.8	9.8	13	15.9	9.1	34	65.2	12.9
Bronx	12	8.7	5.9	27	31.6	11.4	66	59.6	11.7
Kings (Brooklyn)	10	10.7	6.9	22	27.5	11.1	50	61.8	11.9
New York (Manhattan)	9	14.2	10.4	19	23.0	11.1	32	62.8	14.0
Queens	10	13.6	9.7	20	20.8	9.6	60	65.7	11.9
Richmond (Staten Island)	5	9.4	8.3	11	22.3	15.7	26	68.2	16.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**No Health Care Coverage for Past Two Years**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Allegany, Wyoming	25	44.3	33	55.7	14.9
Delaware, Otsego, Schoharie	31	50.9	38	49.1	15.2
Dutchess, Putnam	28	57.4	23	42.6	15.7
Erie	23	57.9	14	42.1	18.7
Fulton, Montgomery	28	47.1	32	52.9	14.4
Niagara	25	68.2	13	31.8	17.6
Onondaga	26	45.5	28	54.5	15.8
Suffolk	22	46.0	32	54.0	16.4
Sullivan, Ulster	35	51.5	47	48.5	13.1
Westchester	27	39.3	32	60.7	14.8
Bronx	43	44.0	60	56.0	11.9
Kings (Brooklyn)	30	35.6	55	64.4	11.9
New York (Manhattan)	24	41.6	33	58.4	16.1
Queens	37	40.0	55	60.0	11.9
Richmond (Staten Island)	19	42.4	25	57.6	17.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Ever Been Tested for HIV, Aged 18-64**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Nassau	187	43.0	255	57.0	5.6
Sullivan, Ulster	254	52.5	232	47.5	5.4

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Ever Been Told Have High Blood Pressure by a Medical Professional**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Albany	161	25.3	466	74.7	3.9
Allegany, Wyoming	180	25.3	434	74.7	3.7
Broome	184	25.8	443	74.2	3.7
Cattaraugus, Chautauqua	234	33.2	384	66.8	4.2
Cayuga, Seneca, Wayne	203	28.2	431	71.8	4.0
Chemung, Schuylar, Tioga	176	26.3	445	73.7	3.8
Chenango, Madison	199	27.3	446	72.7	3.7
Clinton, Essex, Franklin	183	26.7	437	73.3	3.9
Columbia, Greene	182	29.3	439	70.7	4.0
Cortland, Tompkins	133	17.6	497	82.4	3.4
Delaware, Otsego, Schoharie	177	23.8	438	76.2	3.6
Dutchess, Putnam	154	23.6	448	76.4	3.7
Genesee, Orleans	186	28.7	437	71.3	4.0
Hamilton, Herkimer	180	25.5	418	74.5	3.7
Jefferson, Lewis, StLawrence	200	29.3	423	70.7	4.3
Livingston, Ontario	170	25.2	449	74.8	3.8
Nassau	140	22.3	459	77.7	3.8
Niagara	219	31.9	389	68.1	4.0
Oneida, Oswego	181	26.7	461	73.3	3.8
Orange	170	23.2	458	76.8	3.6
Rensselaer	192	28.2	426	71.8	3.9
Rockland	147	22.1	458	77.9	3.6
Saratoga	172	25.9	456	74.1	3.9
Schenectady	189	28.5	433	71.5	3.9
Steuben, Yates	178	25.0	459	75.0	3.7
Suffolk	157	23.2	462	76.8	3.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Had a Fall in Past Three Months**

Locality	n <sup>1</sup>	No fall % <sup>2</sup>	C.I. <sup>3</sup>	n	Fall with injury %	C.I.	n	Fall no injury %	C.I.
Albany	278	79.4	4.9	32	8.7	3.3	41	11.9	4.0
Broome	289	78.5	4.7	29	8.0	3.1	50	13.5	3.9
Cattaraugus, Chautauqua	311	80.5	4.5	23	5.8	2.6	49	13.7	4.0
Cayuga, Seneca, Wayne	315	84.5	3.8	23	4.7	2.1	48	10.8	3.2
Chemung, Schuyler, Tioga	290	79.4	4.6	24	5.9	2.5	52	14.8	4.2
Chenango, Madison	315	81.8	4.0	26	5.8	2.3	52	12.5	3.5
Clinton, Essex, Franklin	287	82.3	4.5	15	4.2	2.4	51	13.5	4.0
Columbia, Greene	318	79.2	4.5	22	3.9	1.7	63	17.0	4.3
Cortland, Tompkins	261	79.1	4.5	28	7.0	2.7	55	13.9	3.8
Delaware, Otsego, Schoharie	320	80.8	4.1	23	5.6	2.4	55	13.6	3.6
Dutchess, Putnam	279	85.9	4.3	15	5.7	3.1	34	8.3	3.2
Genesee, Orleans	290	82.5	4.3	27	6.9	2.8	39	10.6	3.5
Hamilton, Herkimer	314	83.5	4.0	26	7.0	2.8	39	9.4	3.1
Jefferson, Lewis, StLawrence	283	83.3	4.2	25	5.3	2.2	46	11.4	3.7
Livingston, Ontario	297	84.8	4.0	23	5.6	2.6	38	9.6	3.2
Oneida, Oswego	301	82.7	4.4	22	5.4	2.6	42	11.8	3.8
Onondaga	298	84.4	4.1	22	5.9	2.7	36	9.7	3.3
Orange	269	81.5	4.6	27	6.9	2.8	40	11.6	3.9
Rensselaer	283	80.8	4.7	24	5.8	2.8	49	13.4	4.1
Rockland	279	83.1	4.4	24	6.8	3.0	39	10.1	3.5
Saratoga	292	82.4	4.7	18	6.1	3.2	41	11.5	3.7
Schenectady	301	82.4	4.3	20	4.9	2.3	50	12.7	3.8
Steuben, Yates	308	82.8	4.3	21	5.3	2.6	49	11.9	3.6
Suffolk	293	88.7	3.7	17	4.2	2.2	24	7.1	3.1
Westchester	280	87.2	4.1	16	4.3	2.4	30	8.6	3.5
Bronx	237	84.3	5.0	18	7.1	3.6	25	8.6	3.8
Kings (Brooklyn)	218	80.8	5.8	25	7.9	3.4	26	11.3	5.1
New York (Manhattan)	254	88.1	4.0	17	6.0	3.2	21	5.9	2.7
Queens	227	85.6	4.9	8	3.1	2.5	31	11.4	4.3
Richmond (Staten Island)	271	84.6	4.8	11	3.8	2.6	36	11.6	4.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Had a Flu Shot in the Past 12 Months, Aged 65 and Older**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Albany	87	78.3	23	21.7	8.2
Allegany, Wyoming	97	73.5	32	26.5	8.6
Broome	113	65.7	47	34.3	8.6
Cattaraugus, Chautauqua	126	71.8	51	28.2	7.5
Cayuga, Seneca, Wayne	120	71.9	50	28.1	7.4
Chemung, Schuyler, Tioga	102	74.0	43	26.0	7.6
Chenango, Madison	97	69.9	51	30.1	7.9
Clinton, Essex, Franklin	103	69.8	43	30.2	8.3
Columbia, Greene	79	57.8	53	42.2	9.4
Cortland, Tompkins	81	70.4	33	29.6	9.2
Dutchess, Putnam	65	59.0	46	41.0	10.2
Genesee, Orleans	97	72.2	41	27.8	8.0
Hamilton, Herkimer	97	60.7	57	39.3	8.9
Jefferson, Lewis, StLawrence	103	74.4	37	25.6	8.2
Livingston, Ontario	94	80.9	25	19.1	7.6
Monroe	95	75.4	28	24.6	8.8
Nassau	96	72.3	35	27.7	8.7
Niagara	108	70.5	52	29.5	7.4
Oneida, Oswego	89	61.0	58	39.0	8.7
Onondaga	108	79.8	25	20.2	7.9
Orange	79	66.9	38	33.1	9.1
Rensselaer	103	73.6	39	26.4	8.0
Rockland	89	69.4	34	30.6	9.2
Saratoga	87	67.2	42	32.8	9.5
Schenectady	118	70.4	50	29.6	7.5
Steuben, Yates	97	73.3	34	26.7	8.5
Suffolk	84	70.1	39	29.9	8.9
Sullivan, Ulster	93	71.7	39	28.3	8.3
Westchester	97	67.8	39	32.2	9.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health  
**Had a Flu Shot in the Past 12 Months**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Albany	229	35.3	393	64.7	4.2
Allegany, Wyoming	229	33.3	385	66.7	4.1
Broome	204	28.5	420	71.5	3.8
Cattaraugus, Chautauqua	236	32.9	381	67.1	4.1
Cayuga, Seneca, Wayne	235	32.4	400	67.6	4.1
Chemung, Schuyler, Tioga	227	33.7	393	66.3	4.1
Chenango, Madison	226	32.6	418	67.4	4.0
Clinton, Essex, Franklin	221	31.9	397	68.1	4.3
Columbia, Greene	199	30.5	416	69.5	4.0
Cortland, Tompkins	229	31.1	401	68.9	4.1
Dutchess, Putnam	164	25.2	434	74.8	3.9
Genesee, Orleans	229	34.6	394	65.4	4.1
Hamilton, Herkimer	216	32.4	379	67.6	4.2
Jefferson, Lewis, StLawrence	244	36.5	379	63.5	4.6
Livingston, Ontario	259	39.1	356	60.9	4.3
Monroe	236	35.2	373	64.8	4.1
Nassau	216	33.0	383	67.0	4.3
Niagara	210	31.3	400	68.7	4.1
Oneida, Oswego	215	32.2	424	67.8	4.1
Onondaga	243	34.6	370	65.4	4.2
Orange	207	31.6	418	68.4	4.4
Rensselaer	234	35.2	382	64.8	4.2
Rockland	192	28.1	409	71.9	3.9
Saratoga	226	33.0	400	67.0	4.2
Schenectady	254	36.9	369	63.1	4.2
Steuben, Yates	222	32.4	410	67.6	4.1
Suffolk	180	27.9	437	72.1	4.1
Sullivan, Ulster	205	32.2	419	67.8	4.5
Westchester	215	32.0	375	68.0	4.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Ever Had a Pneumonia Shot, Aged 65 and Older**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Albany	80	74.6	26	25.4	8.9
Allegany, Wyoming	92	70.9	36	29.1	8.7
Broome	107	60.6	54	39.4	8.7
Cattaraugus, Chautauqua	122	71.6	50	28.4	7.7
Cayuga, Seneca, Wayne	115	70.4	48	29.6	7.7
Chemung, Schuyler, Tioga	101	75.6	39	24.4	7.6
Chenango, Madison	108	76.9	37	23.1	7.3
Clinton, Essex, Franklin	90	64.5	50	35.5	8.8
Columbia, Greene	81	61.8	49	38.2	9.4
Cortland, Tompkins	85	75.4	27	24.6	8.7
Dutchess, Putnam	74	75.0	33	25.0	8.4
Genesee, Orleans	90	69.3	43	30.7	8.4
Hamilton, Herkimer	104	67.5	46	32.5	8.9
Jefferson, Lewis, StLawrence	106	77.0	30	23.0	8.5
Livingston, Ontario	91	79.8	24	20.2	8.1
Monroe	92	80.0	22	20.0	8.5
Nassau	92	72.1	36	27.9	8.6
Niagara	111	68.3	48	31.7	7.9
Oneida, Oswego	96	66.8	47	33.2	8.5
Onondaga	102	75.4	29	24.6	8.6
Orange	70	60.4	44	39.6	9.7
Rensselaer	96	69.5	45	30.5	8.3
Rockland	76	60.0	44	40.0	9.8
Saratoga	89	75.8	33	24.2	8.9
Schenectady	123	75.0	41	25.0	7.3
Steuben, Yates	87	65.2	43	34.8	9.0
Suffolk	77	60.3	44	39.7	9.9
Sullivan, Ulster	78	60.0	49	40.0	9.6
Westchester	71	53.0	54	47.0	9.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Ever Had a Pneumonia Shot**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Albany	152	25.7	422	74.3	4.0
Allegany, Wyoming	166	25.3	422	74.7	3.9
Broome	173	25.6	418	74.4	3.8
Cattaraugus, Chautauqua	187	28.1	390	71.9	4.1
Cayuga, Seneca, Wayne	171	22.4	426	77.6	3.5
Chemung, Schuyler, Tioga	157	25.2	417	74.8	3.9
Chenango, Madison	192	29.5	411	70.5	4.1
Clinton, Essex, Franklin	172	28.3	403	71.7	4.3
Columbia, Greene	152	25.6	428	74.4	4.0
Cortland, Tompkins	140	21.2	443	78.8	3.9
Dutchess, Putnam	132	23.8	430	76.2	4.1
Genesee, Orleans	148	24.3	435	75.7	3.8
Hamilton, Herkimer	173	26.5	389	73.5	4.0
Jefferson, Lewis, StLawrence	192	30.6	382	69.4	4.6
Livingston, Ontario	169	27.0	407	73.0	4.1
Monroe	144	25.4	412	74.6	4.0
Nassau	141	24.8	420	75.2	4.1
Niagara	178	27.1	396	72.9	3.9
Oneida, Oswego	160	25.1	437	74.9	3.9
Onondaga	187	31.1	382	68.9	4.5
Orange	118	20.0	469	80.0	3.9
Rensselaer	171	28.4	396	71.6	4.1
Rockland	116	18.3	436	81.7	3.4
Saratoga	156	25.8	422	74.2	4.1
Schenectady	192	29.3	393	70.7	4.0
Steuben, Yates	145	23.1	451	76.9	3.8
Suffolk	119	18.0	464	82.0	3.5
Sullivan, Ulster	149	25.3	435	74.7	4.2
Westchester	133	22.2	415	77.8	4.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Have Had Permanent Teeth Removed because of Tooth Decay or Gum Disease**

Locality	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Albany	282	43.5	327	56.5	4.4
Allegany, Wyoming	335	51.2	268	48.8	4.6
Broome	323	48.3	287	51.7	4.5
Cattaraugus, Chautauqua	392	58.3	218	41.7	4.6
Cayuga, Seneca, Wayne	377	54.8	252	45.2	4.6
Chemung, Schuylar, Tioga	366	56.5	250	43.5	4.4
Chenango, Madison	382	55.8	251	44.2	4.5
Clinton, Essex, Franklin	352	52.5	255	47.5	4.7
Columbia, Greene	353	54.9	251	45.1	4.5
Cortland, Tompkins	299	39.3	322	60.7	4.4
Delaware, Otsego, Schoharie	348	52.8	255	47.2	4.8
Fulton, Montgomery	377	58.9	228	41.1	4.5
Genesee, Orleans	338	50.7	275	49.3	4.5
Hamilton, Herkimer	353	54.9	228	45.1	4.8
Jefferson, Lewis, StLawrence	368	52.7	238	47.3	5.0
Livingston, Ontario	302	46.0	305	54.0	4.5
Oneida, Oswego	364	52.7	263	47.3	4.5
Onondaga	314	48.2	293	51.8	4.6
Orange	342	51.0	272	49.0	4.7
Rensselaer	326	51.5	277	48.5	4.5
Rockland	308	48.9	282	51.1	4.6
Saratoga	305	44.6	313	55.4	4.5
Schenectady	339	50.6	270	49.4	4.5
Steuben, Yates	366	56.3	256	43.7	4.6
Warren, Washington	338	49.6	279	50.4	4.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Having Seen a Dental Professional in the Past 12 Months**

Locality	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Albany	468	78.3	127	21.7	3.8
Allegany, Wyoming	374	66.7	172	33.3	4.6
Broome	445	76.1	132	23.9	4.0
Cattaraugus, Chautauqua	379	67.6	159	32.4	4.7
Cayuga, Seneca, Wayne	429	74.1	135	25.9	4.3
Chemung, Schuylar, Tioga	395	70.6	155	29.4	4.3
Chenango, Madison	387	68.7	163	31.3	4.5
Clinton, Essex, Franklin	389	69.2	165	30.8	4.7
Columbia, Greene	417	72.0	151	28.0	4.2
Cortland, Tompkins	414	72.1	155	27.9	4.4
Delaware, Otsego, Schoharie	409	74.2	139	25.8	4.4
Fulton, Montgomery	385	71.6	151	28.4	4.3
Genesee, Orleans	431	75.9	135	24.1	4.1
Hamilton, Herkimer	379	72.3	139	27.7	4.6
Jefferson, Lewis, StLawrence	402	71.4	147	28.6	4.9
Livingston, Ontario	484	83.0	96	17.0	3.5
Oneida, Oswego	452	75.2	136	24.8	4.1
Onondaga	462	78.8	112	21.2	4.2
Orange	427	73.7	161	26.3	4.1
Rensselaer	428	75.6	138	24.4	4.0
Rockland	474	82.2	103	17.8	3.4
Saratoga	469	78.4	114	21.6	4.0
Schenectady	453	77.3	119	22.7	4.0
Steuben, Yates	390	67.7	176	32.3	4.6
Warren, Washington	428	75.5	140	24.5	4.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

**Work-Related Physical Activity**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Mostly sitting or standing %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Mostly walking %</b>	<b>C.I.</b>	<b>n</b>	<b>Mostly heavy labor %</b>	<b>C.I.</b>
Erie	211	64.0	6.0	68	21.9	5.1	42	14.1	4.4
Monroe	238	67.8	5.6	79	25.1	5.3	27	7.1	2.9
Sullivan, Ulster	235	54.8	6.1	73	22.4	5.6	76	22.8	5.0
Warren, Washington	205	55.3	5.9	81	23.4	5.0	68	21.3	5.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

**Vigorous Physical Activity (Vigorous Exercise: 20 or more minutes per day and 3 or more days per week)**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Meet Vig. Objective %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Insufficient Vig. Activity %</b>	<b>C.I.</b>	<b>n</b>	<b>No Vig. Activity %</b>	<b>C.I.</b>
Erie	171	31.0	4.4	96	18.5	3.7	317	50.6	4.6
Monroe	168	28.5	4.1	124	22.2	3.8	308	49.3	4.5
Sullivan, Ulster	166	28.5	4.3	98	18.6	3.9	340	52.9	4.8
Warren, Washington	164	28.1	4.1	126	22.1	3.9	330	49.8	4.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health  
**Meet Recommendations for Moderate or Vigorous Physical Activity**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Meet Objective %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Insufficient Activity %</b>	<b>C.I.</b>	<b>n</b>	<b>No Activity %</b>	<b>C.I.</b>
Erie	278	51.0	4.6	212	36.0	4.4	89	13.0	2.8
Monroe	295	51.0	4.5	239	41.1	4.5	55	7.9	2.3
Sullivan, Ulster	298	50.3	4.9	218	37.7	4.7	71	12.0	3.6
Warren, Washington	326	54.1	4.5	202	34.7	4.4	75	11.1	2.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Any Physical Activity or Exercise**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Erie	555	89.1	77	10.9	2.5
Monroe	573	93.5	53	6.5	1.9
Sullivan, Ulster	572	90.9	71	9.1	2.4
Warren, Washington	583	91.1	65	8.9	2.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Had a Severe Sunburn Lasting 12 Hours in Past 12 Months**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Allegany, Wyoming	249	46.4	364	53.6	4.5
Onondaga	209	38.8	401	61.2	4.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Number of Severe Sunburns Lasting 12 or More Hours in Past 12 Months**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>3 or more %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>1-2 %</b>	<b>C.I.</b>	<b>n</b>	<b>None %</b>	<b>C.I.</b>
Allegany, Wyoming	110	22.7	4.1	128	22.5	3.8	364	54.8	4.6
Onondaga	78	14.6	3.4	128	23.8	4.2	401	61.6	4.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

**Neighborhood Safety**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Extremely or quite safe %<sup>2</sup></b>	<b>n</b>	<b>Slightly or not safe %</b>	<b>C.I.<sup>3</sup></b>
Fulton, Montgomery	551	90.9	59	9.1	2.5
Nassau	558	92.2	32	7.8	3.3
Onondaga	511	81.7	100	18.3	4.0
Suffolk	555	89.2	62	10.8	3.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Living Arrangements**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Own %<sup>2</sup></b>	<b>n</b>	<b>Rent %</b>	<b>C.I.<sup>3</sup></b>
Fulton, Montgomery	436	75.4	170	24.6	3.8
Nassau	473	80.2	117	19.8	3.8
Onondaga	416	70.3	193	29.7	4.3
Suffolk	477	78.5	132	21.5	3.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Lived at Current Address 2 or More Years**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Fulton, Montgomery	496	82.0	120	18.0	3.3
Nassau	522	85.7	71	14.3	3.5
Onondaga	489	79.0	122	21.0	4.0
Suffolk	500	81.4	117	18.6	3.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Concerned about Having Enough Food for You or Your Family in Past 30 Days**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Fulton, Montgomery	41	5.9	573	94.1	2.0
Nassau	27	4.9	567	95.1	2.2
Onondaga	35	4.9	577	95.1	1.8
Suffolk	44	7.5	574	92.5	2.4

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health  
**Number of Cigarettes Smoked per day among Everyday Smokers**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>1-10 cigarettes %<sup>2</sup></b>	<b>n</b>	<b>≥11 cigarettes %</b>	<b>C.I.<sup>3</sup></b>
Allegany, Wyoming	34	26.1	94	73.9	8.9
Delaware, Otsego, Schoharie	35	31.7	75	68.3	10.3
Dutchess, Putnam	35	38.4	68	61.6	12.2
Erie	35	40.5	63	59.5	11.7
Nassau	29	45.6	44	54.4	13.2
Niagara	36	26.6	95	73.4	8.4
Onondaga	41	37.5	71	62.5	10.4
Sullivan, Ulster	32	36.9	73	63.1	11.5
Westchester	34	60.9	30	39.1	13.8
Bronx	51	63.0	32	37.0	12.2
Kings (Brooklyn)	37	49.5	38	50.5	13.1
New York (Manhattan)	40	57.0	31	43.0	13.2
Queens	29	64.0	19	36.0	15.9
Richmond (Staten Island)	38	37.5	63	62.5	11.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Heard, Read or Seen Information about the Effects of Tobacco from Television**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Delaware, Otsego, Schoharie	385	67.6	213	32.4	4.3
Dutchess, Putnam	370	63.8	218	36.2	4.5
Bronx	416	68.8	177	31.2	4.6
Kings (Brooklyn)	379	63.6	221	36.4	4.5
New York (Manhattan)	331	61.2	240	38.8	4.6
Queens	349	63.6	214	36.4	4.7
Richmond (Staten Island)	361	62.2	221	37.8	4.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Heard, Read or Seen Information about the Effects of Tobacco from Radio**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Delaware, Otsego, Schoharie	212	37.1	387	62.9	4.7
Dutchess, Putnam	161	28.5	424	71.5	4.4
Bronx	230	39.2	356	60.8	4.8
Kings (Brooklyn)	192	32.6	407	67.4	4.3
New York (Manhattan)	114	21.1	457	78.9	4.0
Queens	184	33.5	375	66.5	4.6
Richmond (Staten Island)	200	36.0	383	64.0	4.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Heard, Read or Seen Information about the Effects of Tobacco from Billboards**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Delaware, Otsego, Schoharie	189	33.4	418	66.6	4.6
Dutchess, Putnam	225	39.4	362	60.6	4.7
Bronx	278	50.7	312	49.3	4.9
Kings (Brooklyn)	215	39.4	381	60.6	4.7
New York (Manhattan)	195	36.6	376	63.4	4.7
Queens	189	37.7	365	62.3	4.9
Richmond (Staten Island)	218	37.1	365	62.9	4.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Heard, Read or Seen Information about the Effects of Tobacco from Buses**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Delaware, Otsego, Schoharie	24	4.6	581	95.4	2.0
Dutchess, Putnam	33	6.1	556	93.9	2.5
Bronx	218	36.4	371	63.6	4.7
Kings (Brooklyn)	155	27.8	436	72.2	4.3
New York (Manhattan)	153	30.5	413	69.5	4.7
Queens	157	30.0	401	70.0	4.6
Richmond (Staten Island)	147	25.9	434	74.1	4.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Heard, Read or Seen Information about the Effects of Tobacco from Magazines**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Delaware, Otsego, Schoharie	343	57.1	260	42.9	4.7
Dutchess, Putnam	325	55.4	267	44.6	4.6
Bronx	317	51.9	270	48.1	4.9
Kings (Brooklyn)	296	50.8	308	49.2	4.7
New York (Manhattan)	282	49.8	297	50.2	4.8
Queens	290	53.7	273	46.3	4.9
Richmond (Staten Island)	304	49.8	291	50.2	4.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Heard, Read or Seen Information about the Effects of Tobacco from a Website**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Delaware, Otsego, Schoharie	49	9.2	561	90.8	2.9
Dutchess, Putnam	52	10.1	545	89.9	3.2
Bronx	80	15.1	517	84.9	3.5
Kings (Brooklyn)	71	12.2	536	87.8	3.0
New York (Manhattan)	60	11.0	523	89.0	3.0
Queens	70	13.6	495	86.4	3.4
Richmond (Staten Island)	46	8.3	550	91.7	2.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Heard, Read or Seen Information about the Effects of Tobacco from Movie Theater**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Delaware, Otsego, Schoharie	37	7.4	570	92.6	2.8
Dutchess, Putnam	66	14.1	528	85.9	3.8
Bronx	117	22.6	473	77.4	4.3
Kings (Brooklyn)	86	15.0	519	85.0	3.4
New York (Manhattan)	97	17.3	481	82.7	3.6
Queens	69	13.7	496	86.3	3.5
Richmond (Staten Island)	63	10.7	536	89.3	3.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

**Heard or Seen Anti-Tobacco Advertisement that Refers to the NYS Quitline in Past 30 Days**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Delaware, Otsego, Schoharie	215	40.3	359	59.7	4.8
Dutchess, Putnam	208	38.5	361	61.5	4.7
Bronx	301	53.9	275	46.1	5.0
Kings (Brooklyn)	250	43.3	332	56.7	4.7
New York (Manhattan)	221	43.7	329	56.3	4.9
Queens	240	47.7	302	52.3	5.0
Richmond (Staten Island)	220	39.7	349	60.3	4.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Ever Used or Tried Any Smokeless Tobacco Products such as Chewing Tobacco or Snuff**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Bronx	27	4.4	568	95.6	2.1
Kings (Brooklyn)	31	5.1	576	94.9	2.0
New York (Manhattan)	42	7.2	546	92.8	2.4
Queens	20	3.1	550	96.9	1.5
Richmond (Staten Island)	34	7.3	567	92.7	2.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Work Indoors Most of the Time**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Erie	288	86.6	38	13.4	4.5
Nassau	300	85.9	40	14.1	4.6
Sullivan, Ulster	296	74.4	85	25.6	5.3
Westchester	304	80.6	60	19.4	5.1
Bronx	259	77.6	66	22.4	5.6
Kings (Brooklyn)	291	81.4	59	18.6	5.1
New York (Manhattan)	321	89.9	36	10.1	3.6
Queens	264	74.5	71	25.5	6.0
Richmond (Staten Island)	278	76.3	76	23.7	5.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

**Smoking Policy at Place of Work among Indoor Workers**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Not allowed anywhere %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Allowed in some or all places %</b>	<b>C.I.</b>	<b>n</b>	<b>No official policy %</b>	<b>C.I.</b>
Erie	243	82.7	5.3	29	11.7	4.6	14	5.6	3.2
Nassau	253	84.7	4.9	23	9.0	4.2	21	6.3	2.9
Sullivan, Ulster	247	82.1	5.2	28	10.8	4.3	19	7.2	3.4
Westchester	278	91.7	3.6	14	5.2	3.0	10	3.1	2.1
Bronx	210	75.5	7.5	30	17.1	7.0	17	7.4	4.2
Kings (Brooklyn)	239	81.5	5.2	35	13.7	4.6	12	4.8	2.9
New York (Manhattan)	275	86.5	4.4	25	8.3	3.7	19	5.2	2.6
Queens	235	88.6	4.9	13	6.6	4.2	12	4.8	2.8
Richmond (Staten Island)	248	88.0	5.1	21	9.1	4.3	6	2.8	3.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution  
<sup>2</sup>Weighted Percent  
<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Use Seatbelt when Ride or Drive in a Car (Always, Nearly Always or Never Ride in a Car)**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Allegany, Wyoming	566	91.1	46	8.9	2.7
Delaware, Otsego, Schoharie	570	92.3	43	7.7	2.7
Dutchess, Putnam	547	90.0	51	10.0	3.1
Nassau	555	93.7	40	6.3	2.1
Onondaga	581	94.6	32	5.4	2.1
Warren, Washington	590	94.1	40	5.9	2.0
Bronx	518	86.7	73	13.3	3.4
Kings (Brooklyn)	543	88.6	64	11.4	3.0
New York (Manhattan)	521	90.0	65	10.0	2.8
Queens	514	92.0	50	8.0	2.5
Richmond (Staten Island)	535	88.1	65	11.9	3.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

## Expanded BRFSS, 2003 - New York State Department of Health

**Use Seatbelt when Ride or Drive in a Car (Always or Never ride in a Car)**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Allegany, Wyoming	493	78.0	119	22.0	3.9
Delaware, Otsego, Schoharie	494	78.8	119	21.2	4.0
Dutchess, Putnam	495	79.9	103	20.1	4.1
Nassau	494	84.9	101	15.1	3.1
Onondaga	529	85.2	84	14.8	3.4
Warren, Washington	529	83.8	101	16.2	3.3
Bronx	458	74.9	133	25.1	4.3
Kings (Brooklyn)	465	76.1	142	23.9	4.0
New York (Manhattan)	431	73.8	155	26.2	4.1
Queens	443	78.1	121	21.9	4.1
Richmond (Staten Island)	452	74.5	148	25.5	4.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Trying to Lose Weight**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Albany	290	44.8	333	55.2	4.4
Allegany, Wyoming	259	40.7	354	59.3	4.4
Broome	282	42.9	344	57.1	4.3
Cattaraugus, Chautauqua	292	47.8	323	52.2	4.6
Cayuga, Seneca, Wayne	317	48.8	319	51.2	4.5
Chemung, Schuylar, Tioga	294	46.4	327	53.6	4.4
Chenango, Madison	289	42.7	354	57.3	4.4
Clinton, Essex, Franklin	277	43.5	341	56.5	4.6
Columbia, Greene	299	46.8	319	53.2	4.4
Cortland, Tompkins	275	40.2	355	59.8	4.6
Fulton, Montgomery	291	46.3	324	53.7	4.5
Genesee, Orleans	315	50.7	307	49.3	4.4
Hamilton, Herkimer	279	44.1	317	55.9	4.6
Jefferson, Lewis, StLawrence	283	46.7	341	53.3	4.9
Livingston, Ontario	312	47.3	304	52.7	4.5
Nassau	309	49.4	282	50.6	4.8
Niagara	284	45.7	326	54.3	4.5
Oneida, Oswego	299	44.8	341	55.2	4.4
Onondaga	302	48.3	311	51.7	4.6
Orange	329	50.0	294	50.0	4.7
Rensselaer	316	47.0	298	53.0	4.4
Rockland	316	50.4	288	49.6	4.6
Saratoga	322	50.4	304	49.6	4.6
Schenectady	294	47.2	330	52.8	4.5
Steuben, Yates	310	45.2	324	54.8	4.5
Suffolk	312	49.7	305	50.3	4.5
Westchester	276	45.4	314	54.6	4.7
Bronx	291	45.7	300	54.3	4.9
Kings (Brooklyn)	315	48.7	293	51.3	4.6
New York (Manhattan)	268	44.5	317	55.5	4.7
Queens	283	48.4	281	51.6	4.9
Richmond (Staten Island)	312	50.9	286	49.1	4.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Trying to Maintain Current Weight of Those not Trying to Lose Weight**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Albany	228	67.2	105	32.8	5.8
Allegany, Wyoming	198	53.7	152	46.3	6.0
Broome	206	59.1	133	40.9	6.0
Cattaraugus, Chautauqua	212	66.0	110	34.0	6.0
Cayuga, Seneca, Wayne	199	62.5	118	37.5	6.5
Chemung, Schuylar, Tioga	206	64.1	120	35.9	5.8
Chenango, Madison	213	60.4	142	39.6	5.9
Clinton, Essex, Franklin	193	55.9	145	44.1	6.3
Columbia, Greene	174	53.9	143	46.1	6.2
Cortland, Tompkins	214	58.4	139	41.6	6.1
Fulton, Montgomery	190	60.1	132	39.9	6.1
Genesee, Orleans	192	62.5	115	37.5	6.1
Hamilton, Herkimer	176	53.7	138	46.3	6.6
Jefferson, Lewis, StLawrence	203	61.5	135	38.5	6.4
Livingston, Ontario	189	60.1	113	39.9	6.2
Nassau	177	59.0	107	41.0	6.7
Niagara	226	69.4	96	30.6	5.9
Oneida, Oswego	213	61.8	125	38.2	6.0
Onondaga	181	55.4	127	44.6	6.7
Orange	171	54.7	122	45.3	6.9
Rensselaer	186	61.1	113	38.9	6.3
Rockland	170	60.3	113	39.7	6.6
Saratoga	191	60.2	113	39.8	6.5
Schenectady	203	60.9	125	39.1	6.0
Steuben, Yates	196	57.5	128	42.5	6.6
Suffolk	201	66.5	104	33.5	6.0
Westchester	214	65.9	98	34.1	6.3
Bronx	187	62.9	113	37.1	6.7
Kings (Brooklyn)	171	57.8	120	42.2	6.7
New York (Manhattan)	205	61.3	113	38.7	6.4
Queens	163	58.5	119	41.5	6.7
Richmond (Staten Island)	172	56.7	114	43.3	6.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health  
**Eating Fewer Calories or Less Fat to Lose or Maintain Weight**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Albany	396	75.8	117	24.2	4.3
Allegany, Wyoming	332	72.0	120	28.0	4.8
Broome	379	78.3	106	21.7	4.1
Cattaraugus, Chautauqua	382	73.7	117	26.3	4.7
Cayuga, Seneca, Wayne	384	74.5	123	25.5	4.5
Chemung, Schuylar, Tioga	364	71.9	130	28.1	4.5
Chenango, Madison	386	76.3	110	23.7	4.4
Clinton, Essex, Franklin	354	76.3	110	23.7	4.6
Columbia, Greene	368	78.9	100	21.1	4.2
Cortland, Tompkins	353	68.6	134	31.4	5.2
Fulton, Montgomery	373	79.6	98	20.4	4.1
Genesee, Orleans	392	78.0	111	22.0	4.1
Hamilton, Herkimer	351	76.1	99	23.9	4.7
Jefferson, Lewis, StLawrence	354	74.2	125	25.8	4.8
Livingston, Ontario	391	77.8	104	22.2	4.2
Nassau	374	78.0	105	22.0	4.3
Niagara	387	76.4	117	23.6	4.1
Oneida, Oswego	385	74.8	121	25.2	4.5
Onondaga	367	75.5	105	24.5	4.5
Orange	396	79.0	97	21.0	4.4
Rensselaer	402	81.5	91	18.5	3.8
Rockland	383	76.5	97	23.5	4.7
Saratoga	379	73.7	130	26.3	4.5
Schenectady	378	76.7	110	23.3	4.4
Steuben, Yates	375	73.6	123	26.4	4.5
Suffolk	411	81.4	95	18.6	3.9
Westchester	366	76.1	113	23.9	4.6
Bronx	370	77.3	102	22.7	4.7
Kings (Brooklyn)	368	80.2	105	19.8	3.9
New York (Manhattan)	325	69.1	140	30.9	5.0
Queens	351	79.6	84	20.4	4.9
Richmond (Staten Island)	377	79.3	100	20.7	4.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Using Physical Exercise to Lose or Maintain Weight**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Albany	401	76.8	115	23.2	4.1
Allegany, Wyoming	311	71.2	142	28.8	4.6
Broome	373	77.8	114	22.2	4.1
Cattaraugus, Chautauqua	346	72.3	158	27.7	4.3
Cayuga, Seneca, Wayne	354	72.0	162	28.0	4.3
Chemung, Schuyler, Tioga	344	71.2	155	28.8	4.4
Chenango, Madison	345	69.6	157	30.4	4.5
Clinton, Essex, Franklin	327	72.5	140	27.5	4.6
Columbia, Greene	350	75.7	122	24.3	4.3
Cortland, Tompkins	361	76.3	126	23.7	4.4
Fulton, Montgomery	316	67.2	165	32.8	4.7
Genesee, Orleans	344	71.2	161	28.8	4.3
Hamilton, Herkimer	317	71.8	137	28.2	4.6
Jefferson, Lewis, StLawrence	366	78.1	119	21.9	4.3
Livingston, Ontario	373	77.3	126	22.7	4.0
Nassau	340	69.0	146	31.0	4.9
Niagara	340	71.0	169	29.0	4.2
Oneida, Oswego	366	73.6	144	26.4	4.3
Onondaga	359	76.2	122	23.8	4.3
Orange	358	74.0	141	26.0	4.4
Rensselaer	379	76.8	122	23.2	4.1
Rockland	325	68.8	159	31.2	4.6
Saratoga	374	73.3	139	26.7	4.4
Schenectady	361	75.2	136	24.8	4.1
Steuben, Yates	355	71.7	150	28.3	4.4
Suffolk	380	75.1	131	24.9	4.3
Westchester	348	72.6	140	27.4	4.6
Bronx	295	60.7	182	39.3	5.4
Kings (Brooklyn)	328	68.8	157	31.2	4.7
New York (Manhattan)	337	70.1	136	29.9	4.9
Queens	290	65.9	156	34.1	5.2
Richmond (Staten Island)	321	68.5	160	31.5	4.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Ever Had a Pap Smear**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>
Albany	305	95.9	10	4.1	3.3
Broome	263	95.8	11	4.2	2.7
Cattaraugus, Chautauqua	266	93.3	19	6.7	3.6
Cayuga, Seneca, Wayne	274	94.5	16	5.5	3.5
Chemung, Schuyler, Tioga	289	97.6	8	2.4	1.8
Chenango, Madison	284	96.6	10	3.4	2.5
Clinton, Essex, Franklin	272	97.0	6	3.0	2.6
Columbia, Greene	268	94.9	12	5.1	3.4
Cortland, Tompkins	273	89.4	20	10.6	5.1
Erie	265	97.5	7	2.5	2.0
Genesee, Orleans	290	98.6	5	1.4	1.2
Hamilton, Herkimer	250	95.1	14	4.9	2.9
Jefferson, Lewis, StLawrence	273	97.7	9	2.3	1.7
Livingston, Ontario	288	96.8	6	3.2	2.9
Niagara	266	95.4	9	4.6	3.9
Oneida, Oswego	309	96.5	11	3.5	2.4
Onondaga	309	97.2	8	2.8	2.2
Orange	321	96.4	10	3.6	2.3
Rensselaer	288	95.2	10	4.8	3.3
Rockland	295	92.5	19	7.5	3.6
Saratoga	321	97.5	5	2.5	3.0
Schenectady	293	96.0	10	4.0	2.9
Steuben, Yates	292	96.9	8	3.1	2.3
Suffolk	312	96.4	9	3.6	2.7
Sullivan, Ulster	292	91.9	15	8.1	4.6
Warren, Washington	283	98.5	3	1.5	1.7
Westchester	298	95.5	11	4.5	3.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health  
**Had a Pap Smear in Past 3 Years**

<b>Locality</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Albany	291	92.2	22	7.8	3.9
Broome	240	89.6	34	10.4	3.7
Cattaraugus, Chautauqua	223	79.2	60	20.8	5.5
Cayuga, Seneca, Wayne	232	81.8	54	18.2	5.4
Chemung, Schuyler, Tioga	256	87.7	41	12.3	3.9
Chenango, Madison	251	85.0	41	15.0	4.9
Clinton, Essex, Franklin	240	86.5	38	13.5	4.4
Columbia, Greene	225	81.4	53	18.6	5.1
Cortland, Tompkins	244	81.9	48	18.1	5.7
Erie	240	90.0	31	10.0	3.7
Genesee, Orleans	257	89.0	36	11.0	3.6
Hamilton, Herkimer	217	84.8	45	15.2	4.6
Jefferson, Lewis, StLawrence	241	87.7	39	12.3	4.5
Livingston, Ontario	263	89.7	30	10.3	4.1
Niagara	235	87.2	35	12.8	5.0
Oneida, Oswego	267	84.8	53	15.2	4.2
Onondaga	291	92.4	24	7.6	3.3
Orange	280	85.9	47	14.1	4.1
Rensselaer	265	88.1	32	11.9	4.3
Rockland	273	87.3	40	12.7	4.2
Saratoga	297	92.2	28	7.8	3.7
Schenectady	268	87.6	34	12.4	4.6
Steuben, Yates	274	92.0	24	8.0	3.4
Suffolk	278	87.3	41	12.7	4.1
Sullivan, Ulster	248	80.3	57	19.7	5.6
Warren, Washington	256	90.8	28	9.2	3.7
Westchester	269	86.9	38	13.1	4.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval