

Expanded BRFSS, 2003

Niagara Locality Tables

TABLE OF CONTENTS:

Core 1: Health Status / Healthy Days 1
Core 2: Health Care Access 4
Core 3: Exercise 8
Core 4: Diabetes 9
Core 5: Asthma 10
Core 6: Arthritis 12
Core 7: Tobacco Use 14
Core 8: Tobacco ETS 17
Core 9: Alcohol Consumption 19
Core 10: Weight Status 21
Core 11: Mammography 23
Core 12: Sexual Behavior 25
Core 13: Family Planning 28
Core 14: Cardiovascular Disease 29
Core 15: Prostate Cancer Screening 31
Core 16: Colorectal Cancer Screening 37

New York State Department Of Health

Expanded BRFSS, 2003 - New York State Department of Health

Self-rated Health Status

Niagara

Demographic Groups	n¹	Fair or Poor %²	n	Good to Excellent %	C.I.³
Total	122	16.4	507	83.6	3.0
Gender					
Male	48	16.7	216	83.3	4.7
Female	74	16.2	291	83.8	3.8
Age					
18-34	8	5.9	101	94.1	4.3
35-54	36	12.7	223	87.3	4.4
≥55	78	29.7	177	70.3	6.0
Education					
≤High School	71	21.5	208	78.5	5.0
>High School	50	12.2	298	87.8	3.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Reported Number of Physical Distress Days in the Past Month

Niagara

Demographic Groups	n ¹	≥14 days % ²	n	< 14 days %	C.I. ³
Total	89	12.3	525	87.7	2.7
Gender					
Male	35	13.1	225	86.9	4.4
Female	54	11.6	300	88.4	3.3
Age					
18-34	6	4.7	103	95.3	4.0
35-54	38	13.3	220	86.7	4.5
≥55	45	17.9	196	82.1	5.2
Education					
≤High School	50	14.9	222	85.1	4.3
>High School	39	10.2	303	89.8	3.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Reported Number of Mental Distress Days in the Past Month

Niagara

Demographic Groups	n ¹	≥14 days % ²	n	< 14 days %	C.I. ³
Total	69	9.8	559	90.2	2.5
Gender					
Male	24	9.0	241	91.0	3.8
Female	45	10.6	318	89.4	3.4
Age					
18-34	11	10.9	98	89.1	6.4
35-54	36	10.6	222	89.4	3.6
≥55	22	8.2	233	91.8	3.6
Education					
≤High School	31	10.7	247	89.3	4.1
>High School	38	9.1	311	90.9	3.1

¹ Percents based on row denominators of less than 50 are unstable and should be used with caution

² Weighted Percent

³ 95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Health Insurance Coverage, 18-64 Years of Age

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	423	88.9	40	11.1	3.8
Gender					
Male	190	87.6	18	12.4	5.9
Female	233	90.1	22	9.9	4.9
Age					
18-34	91	78.0	18	22.0	9.5
35-54	243	93.8	19	6.2	2.9
55-64	89	96.5	3	3.5	4.1
Education					
≤High School	155	82.7	25	17.3	7.1
>High School	267	93.2	15	6.8	4.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Health Insurance Coverage, Aged 18 and Older

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	594	91.2	40	8.8	3.1
Gender					
Male	249	89.9	18	10.1	4.9
Female	345	92.4	22	7.6	3.8
Age					
18-34	91	78.0	18	22.0	9.5
35-54	243	93.8	19	6.2	2.9
≥55	254	98.6	3	1.4	1.6
Education					
≤High School	257	87.4	25	12.6	5.4
>High School	335	94.2	15	5.8	3.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Reported Having a Routine Physical Exam in Past Two Years

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	579	90.7	52	9.3	2.9
Gender					
Male	236	88.4	29	11.6	4.5
Female	343	92.7	23	7.3	3.6
Age					
18-34	92	83.9	16	16.1	8.2
35-54	234	89.8	27	10.2	4.0
≥55	249	97.2	8	2.8	2.1
Education					
≤High School	262	91.7	20	8.3	3.9
>High School	315	89.8	32	10.2	4.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Time in Past 12 Months when Needed Medical Care but Could Not Get It

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	26	4.3	606	95.7	2.0
Gender					
Male	11	4.3	256	95.7	2.6
Female	15	4.3	350	95.7	2.9
Age					
18-34	7	6.6	102	93.4	5.7
35-54	14	4.9	248	95.1	2.7
≥55	5	2.0	250	98.0	1.8
Education					
≤High School	14	5.6	268	94.4	3.6
>High School	12	3.3	336	96.7	2.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	462	74.5	172	25.5	3.8
Gender					
Male	204	77.7	63	22.3	5.5
Female	258	71.6	109	28.4	5.3
Age					
18-34	85	75.4	24	24.6	9.3
35-54	209	82.1	53	17.9	4.8
≥55	162	64.2	95	35.8	6.3
Education					
≤High School	188	69.6	94	30.4	5.9
>High School	274	78.8	76	21.2	5.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Physician Diagnosed Diabetes

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	59	7.7	574	92.3	2.1
Gender					
Male	19	6.8	247	93.2	3.2
Female	40	8.5	327	91.5	2.9
Age					
18-34	1	0.9	108	99.1	1.8
35-54	15	5.7	247	94.3	3.2
≥55	43	15.7	213	84.3	4.7
Education					
≤High School	28	8.0	254	92.0	3.2
>High School	31	7.5	318	92.5	2.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Lifetime Asthma Diagnosed by a Medical Professional

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	82	12.3	551	87.7	2.8
Gender					
Male	26	9.2	241	90.8	3.6
Female	56	15.2	310	84.8	4.2
Age					
18-34	19	14.1	90	85.9	6.6
35-54	25	9.6	237	90.4	4.0
≥55	38	14.4	218	85.6	4.6
Education					
≤High School	43	14.3	239	85.7	4.5
>High School	39	10.7	310	89.3	3.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Current Asthma Diagnosed by a Medical Professional

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	61	9.4	567	90.6	2.6
Gender					
Male	18	6.6	246	93.4	3.1
Female	43	12.0	321	88.0	4.0
Age					
18-34	15	11.3	93	88.7	6.1
35-54	20	7.6	241	92.4	3.7
≥55	26	10.2	227	89.8	4.0
Education					
≤High School	34	11.9	246	88.1	4.2
>High School	27	7.5	319	92.5	3.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Arthritis Diagnosed by a Medical Professional

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	233	32.6	401	67.4	3.9
Gender					
Male	86	29.1	181	70.9	5.7
Female	147	35.7	220	64.3	5.4
Age					
18-34	15	12.6	94	87.4	6.3
35-54	68	23.4	194	76.6	5.4
≥55	150	60.2	107	39.8	6.4
Education					
≤High School	127	39.8	155	60.2	6.3
>High School	104	26.4	246	73.6	5.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

Niagara

Demographic Groups	n ¹	Doctor Diagnosed % ²	C.I. ³	n	Possible Arthritis (CJS Only) %	C.I.	n	No Arthritis or CJS %	C.I.
Total	233	32.8	4.0	103	17.6	3.4	294	49.6	4.4
Gender									
Male	86	29.3	5.8	51	20.0	5.4	128	50.7	6.7
Female	147	36.0	5.4	52	15.4	4.3	166	48.6	5.9
Age									
18-34	15	12.7	6.4	17	16.4	7.6	76	70.9	9.2
35-54	68	23.6	5.5	54	22.5	5.8	139	53.9	6.6
≥55	150	60.4	6.4	31	12.6	4.5	75	27.0	5.7
Education									
≤High School	127	39.8	6.3	40	17.4	5.4	114	42.8	6.7
>High School	104	26.8	5.0	63	17.8	4.4	180	55.4	5.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Smoked 100 Cigarettes in Lifetime

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	376	57.0	257	43.0	4.4
Gender					
Male	176	61.9	90	38.1	6.7
Female	200	52.5	167	47.5	5.9
Age					
18-34	56	49.6	53	50.4	10.4
35-54	156	57.1	106	42.9	6.6
≥55	162	63.2	94	36.8	6.4
Education					
≤High School	184	64.6	98	35.4	6.4
>High School	191	50.7	158	49.3	5.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Current Smoker

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	173	26.6	460	73.4	3.8
Gender					
Male	77	26.6	189	73.4	5.7
Female	96	26.6	271	73.4	5.1
Age					
18-34	42	35.5	67	64.5	9.6
35-54	86	29.5	176	70.5	5.8
≥55	44	16.1	212	83.9	4.7
Education					
≤High School	92	32.8	190	67.2	6.2
>High School	80	21.4	269	78.6	4.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Status of Cigarette Smoking*Niagara*

Demographic Groups	n¹	Everyday Smoker %²	C.I.³	n	Someday Smoker %	C.I.	n	Former or Never Smoked %	C.I.
Total	142	22.2	3.6	31	4.4	1.7	460	73.4	3.8
Gender									
Male	64	23.2	5.5	13	3.4	2.0	189	73.4	5.7
Female	78	21.3	4.7	18	5.2	2.7	271	73.4	5.1
Age									
18-34	36	29.7	9.1	6	5.8	4.6	67	64.5	9.6
35-54	73	25.2	5.5	13	4.3	2.4	176	70.5	5.8
≥55	32	12.6	4.3	12	3.5	2.0	212	83.9	4.7
Education									
≤High School	78	28.9	6.0	14	3.9	2.3	190	67.2	6.2
>High School	63	16.5	4.1	17	4.8	2.5	269	78.6	4.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution²Weighted Percent³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Home Smoking Policies*Niagara*

Demographic Groups	n¹	Not Allowed %²	C.I.³	n	Allowed Some Places %	C.I.	n	Allowed Anywhere %	C.I.
Total	361	59.1	4.3	106	16.9	3.3	166	24.0	3.7
Gender									
Male	141	55.5	6.6	37	14.0	4.6	88	30.6	6.1
Female	220	62.3	5.6	69	19.6	4.7	78	18.1	4.1
Age									
18-34	56	52.8	10.3	29	26.2	8.9	24	21.0	8.3
35-54	148	60.6	6.3	53	17.3	4.7	61	22.1	5.4
≥55	153	62.4	6.3	23	8.8	3.6	80	28.8	5.9
Education									
≤High School	133	49.3	6.7	52	19.7	5.4	96	31.1	6.0
>High School	227	67.1	5.5	53	14.5	4.1	70	18.4	4.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution²Weighted Percent³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Home Smoking

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	201	31.2	433	68.8	4.0
Gender					
Male	88	32.2	179	67.8	6.2
Female	113	30.3	254	69.7	5.3
Age					
18-34	45	39.3	64	60.7	10.0
35-54	92	31.5	170	68.5	6.0
≥55	63	24.5	194	75.5	5.6
Education					
≤High School	111	41.3	171	58.7	6.6
>High School	89	22.8	261	77.2	4.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	83	15.8	544	84.2	3.5
Gender					
Male	59	24.2	203	75.8	5.9
Female	24	8.4	341	91.6	3.9
Age					
18-34	30	29.2	79	70.8	9.6
35-54	42	16.9	215	83.1	5.1
≥55	11	4.0	246	96.0	2.4
Education					
≤High School	43	19.7	237	80.3	5.9
>High School	40	12.7	305	87.3	4.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	40	6.5	588	93.5	2.2
Gender					
Male	22	8.8	241	91.2	3.9
Female	18	4.5	347	95.5	2.3
Age					
18-34	10	9.0	99	91.0	5.8
35-54	19	7.1	240	92.9	3.4
≥55	11	3.9	245	96.1	2.4
Education					
≤High School	21	8.3	258	91.7	3.8
>High School	19	5.1	328	94.9	2.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Overweight or Obese, BMI(Body Mass Index) 25 or More

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	363	60.1	240	39.9	4.4
Gender					
Male	189	70.9	76	29.1	6.1
Female	174	49.7	164	50.3	6.1
Age					
18-34	53	49.2	54	50.8	10.5
35-54	150	61.7	100	38.3	6.5
≥55	158	67.0	85	33.0	6.3
Education					
≤High School	168	64.1	96	35.9	6.6
>High School	194	56.8	143	43.2	6.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+)

Niagara

Demographic Groups	n¹	Underweight\ Normal %²	C.I.³	n	Overweight %	C.I.	n	Obese %	C.I.
Total	240	39.9	4.4	222	36.2	4.3	141	23.9	3.9
Gender									
Male	76	29.1	6.1	117	43.0	6.6	72	27.9	6.1
Female	164	50.3	6.1	105	29.7	5.5	69	20.0	4.9
Age									
18-34	54	50.8	10.5	27	25.6	9.2	26	23.6	8.9
35-54	100	38.3	6.5	93	37.7	6.5	57	24.1	6.1
≥55	85	33.0	6.3	101	43.4	6.7	57	23.6	5.8
Education									
≤High School	96	35.9	6.6	99	35.5	6.5	69	28.6	6.3
>High School	143	43.2	6.0	123	36.9	5.8	71	19.9	4.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had Mammography Screening, Women Aged 40 Years and Older

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Female	233	90.2	30	9.8	3.7
Age					
40-64	137	90.2	18	9.8	4.9
≥65	96	90.4	12	9.6	5.5
Education					
≤High School	111	89.8	14	10.2	5.7
>High School	121	90.6	16	9.4	4.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Female	199	78.3	62	21.7	5.3
Age					
40-64	121	81.2	33	18.8	6.4
≥65	78	73.1	29	26.9	8.9
Education					
≤High School	97	78.7	28	21.3	7.6
>High School	101	77.9	34	22.1	7.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health
Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	49	12.7	405	87.3	3.9
Gender					
Male	17	9.7	187	90.3	5.1
Female	32	15.6	218	84.4	5.8
Age					
18-34	30	29.6	78	70.4	9.7
35-54	17	4.4	238	95.6	2.3
55-64	2	2.5	89	97.5	3.5
Education					
≤High School	22	15.8	157	84.2	7.1
>High School	27	10.6	247	89.4	4.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Sexually Active, Age 18-64 Years Old

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	349	84.7	75	15.3	3.9
Gender					
Male	162	84.0	32	16.0	5.9
Female	187	85.5	43	14.5	5.0
Age					
18-34	94	87.0	11	13.0	8.0
35-54	201	88.0	35	12.0	4.2
55-64	54	70.0	29	30.0	10.2
Education					
≤High School	138	84.7	28	15.3	6.0
>High School	210	84.7	47	15.3	5.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	58	17.3	301	82.7	4.8
Gender					
Male	35	19.4	131	80.6	6.8
Female	23	15.3	170	84.7	6.8
Age					
18-34	28	31.9	66	68.1	10.9
35-54	26	10.1	183	89.9	4.0
55-64	4	5.5	52	94.5	5.5
Education					
≤High School	27	22.9	116	77.1	8.8
>High School	31	13.3	184	86.7	5.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)

Niagara

Demographic Groups	Use BC			No BC at risk			No BC		
	n ¹	% ²	C.I. ³	n	%	C.I.	n	%	C.I.
Total	216	67.7	5.8	39	12.8	4.0	65	19.5	5.0
Gender									
Male	115	62.0	7.8	24	13.5	5.3	44	24.4	7.1
Female	101	75.5	8.2	15	11.7	6.0	21	12.7	6.5
Age									
18-34	68	63.4	10.1	17	15.0	7.0	22	21.7	9.0
35-59	148	71.1	6.4	22	11.1	4.5	43	17.8	5.3
Education									
≤High School	88	70.8	9.0	14	11.2	5.9	22	17.9	7.8
>High School	128	65.9	7.5	25	13.9	5.4	42	20.2	6.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Physician Diagnosed Heart Attack, Angina or Stroke

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	73	9.8	550	90.2	2.4
Gender					
Male	34	11.9	227	88.1	4.0
Female	39	8.1	323	91.9	2.7
Age					
18-64	26	4.8	428	95.2	2.0
≥65	47	29.6	118	70.4	7.6
Education					
≤High School	44	13.3	234	86.7	4.1
>High School	27	6.6	316	93.4	2.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina)

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	65	8.8	558	91.2	2.3
Gender					
Male	32	11.1	229	88.9	3.9
Female	33	6.8	329	93.2	2.5
Age					
18-64	23	4.2	431	95.8	1.8
≥65	42	26.8	123	73.2	7.5
Education					
≤High School	40	11.9	238	88.1	3.9
>High School	23	5.9	320	94.1	2.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Male	121	67.0	55	33.0	7.6
Age					
40-64	76	59.4	48	40.6	9.3
≥65	45	86.4	7	13.6	10.3
Education					
≤High School	47	59.2	27	40.8	12.2
>High School	74	72.6	28	27.4	9.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Male	111	61.7	63	38.3	7.8
Age					
40-64	68	53.2	54	46.8	9.5
≥65	43	83.0	9	17.0	11.1
Education					
≤High School	44	54.9	30	45.1	12.2
>High School	67	66.7	33	33.3	9.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had a Digital Rectal Exam, Men Aged 40 and Older

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Male	153	81.8	32	18.2	6.1
Age					
40-64	102	77.7	28	22.3	7.8
≥65	51	92.5	4	7.5	8.1
Education					
≤High School	62	75.5	18	24.5	10.6
>High School	91	87.3	13	12.7	6.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Used a Blood Stool Test at Home, Aged 50 and Older

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	151	49.7	157	50.3	6.1
Gender					
Male	59	49.3	61	50.7	9.6
Female	92	50.1	96	49.9	7.7
Age					
50-64	68	47.2	75	52.8	8.8
≥65	83	52.3	82	47.7	8.2
Education					
≤High School	75	48.9	81	51.1	8.5
>High School	76	50.6	75	49.4	8.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	71	24.9	228	75.1	5.4
Gender					
Male	32	27.9	87	72.1	8.7
Female	39	22.4	141	77.6	6.7
Age					
50-64	33	24.1	109	75.9	7.6
≥65	38	25.9	119	74.1	7.6
Education					
≤High School	36	24.4	115	75.6	7.4
>High School	35	25.6	112	74.4	7.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	98	33.9	201	66.1	5.8
Gender					
Male	42	36.1	77	63.9	9.2
Female	56	32.1	124	67.9	7.4
Age					
50-64	46	33.1	96	66.9	8.3
≥65	52	34.8	105	65.2	8.1
Education					
≤High School	45	30.5	106	69.5	7.9
>High School	53	37.6	94	62.4	8.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older

Niagara

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	160	53.1	144	46.9	6.1
Gender					
Male	71	60.5	46	39.5	9.5
Female	89	47.3	98	52.7	7.7
Age					
50-64	77	54.2	66	45.8	8.8
≥65	83	52.0	78	48.0	8.3
Education					
≤High School	75	49.4	78	50.6	8.6
>High School	84	56.7	66	43.3	8.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	151	50.8	150	49.2	6.1
Gender					
Male	67	57.4	49	42.6	9.7
Female	84	45.6	101	54.4	7.7
Age					
50-64	73	51.7	69	48.3	8.9
≥65	78	50.0	81	50.0	8.4
Education					
≤High School	71	47.0	81	53.0	8.6
>High School	80	54.7	69	45.3	8.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older

Niagara

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	182	60.8	125	39.2	5.9
Gender					
Male	80	67.3	40	32.7	9.0
Female	102	55.5	85	44.5	7.6
Age					
50-64	86	60.7	57	39.3	8.6
≥65	96	60.8	68	39.2	8.0
Education					
≤High School	88	57.6	67	42.4	8.4
>High School	94	64.2	57	35.8	8.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval