

Expanded BRFSS, 2003

New York Locality Tables

TABLE OF CONTENTS:

| | |
|---|----|
| Core 1: Health Status / Healthy Days..... | 1 |
| Core 2: Health Care Access..... | 4 |
| Core 3: Exercise..... | 8 |
| Core 4: Diabetes..... | 9 |
| Core 5: Asthma..... | 10 |
| Core 6: Arthritis..... | 12 |
| Core 7: Tobacco Use..... | 14 |
| Core 8: Tobacco ETS..... | 17 |
| Core 9: Alcohol Consumption..... | 19 |
| Core 10: Weight Status..... | 21 |
| Core 11: Mammography..... | 23 |
| Core 12: Sexual Behavior..... | 25 |
| Core 13: Family Planning..... | 28 |
| Core 14: Cardiovascular Disease..... | 29 |
| Core 15: Prostate Cancer Screening..... | 31 |
| Core 16: Colorectal Cancer Screening..... | 37 |

New York State Department Of Health

Expanded BRFSS, 2003 - New York State Department of Health

Self-rated Health Status

New York

| Demographic Groups | n ¹ | Fair or Poor % ² | n | Good to Excellent % | C.I. ³ |
|--------------------|----------------|-----------------------------|-----|---------------------|-------------------|
| Total | 113 | 18.8 | 501 | 81.2 | 3.9 |
| Gender | | | | | |
| Male | 36 | 16.5 | 209 | 83.5 | 6.3 |
| Female | 77 | 20.7 | 292 | 79.3 | 4.8 |
| Age | | | | | |
| 18-34 | 15 | 12.0 | 166 | 88.0 | 7.2 |
| 35-54 | 29 | 12.8 | 203 | 87.2 | 4.8 |
| ≥55 | 65 | 35.5 | 122 | 64.5 | 8.1 |
| Education | | | | | |
| ≤High School | 58 | 39.7 | 89 | 60.3 | 9.8 |
| >High School | 52 | 9.7 | 411 | 90.3 | 2.8 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Reported Number of Physical Distress Days in the Past Month

New York

| Demographic Groups | n ¹ | ≥14 days % ² | n | < 14 days % | C.I. ³ |
|--------------------|----------------|-------------------------|-----|-------------|-------------------|
| Total | 65 | 8.8 | 533 | 91.2 | 2.3 |
| Gender | | | | | |
| Male | 19 | 6.2 | 222 | 93.8 | 3.0 |
| Female | 46 | 11.1 | 311 | 88.9 | 3.5 |
| Age | | | | | |
| 18-34 | 6 | 2.1 | 173 | 97.9 | 1.8 |
| 35-54 | 21 | 7.8 | 207 | 92.2 | 3.6 |
| ≥55 | 35 | 19.2 | 143 | 80.8 | 6.6 |
| Education | | | | | |
| ≤High School | 25 | 13.5 | 111 | 86.5 | 5.9 |
| >High School | 38 | 6.7 | 420 | 93.3 | 2.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Reported Number of Mental Distress Days in the Past Month

New York

| Demographic Groups | n ¹ | ≥14 days % ² | n | < 14 days % | C.I. ³ |
|--------------------|----------------|-------------------------|-----|-------------|-------------------|
| Total | 78 | 13.2 | 523 | 86.8 | 3.1 |
| Gender | | | | | |
| Male | 21 | 10.7 | 219 | 89.3 | 4.8 |
| Female | 57 | 15.3 | 304 | 84.7 | 4.1 |
| Age | | | | | |
| 18-34 | 29 | 16.2 | 150 | 83.8 | 6.1 |
| 35-54 | 32 | 13.3 | 196 | 86.7 | 4.9 |
| ≥55 | 15 | 8.8 | 167 | 91.2 | 4.9 |
| Education | | | | | |
| ≤High School | 23 | 14.8 | 116 | 85.2 | 6.6 |
| >High School | 54 | 12.5 | 404 | 87.5 | 3.5 |

¹ Percents based on row denominators of less than 50 are unstable and should be used with caution

² Weighted Percent

³ 95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Health Insurance Coverage, 18-64 Years of Age

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|----|------|-------------------|
| Total | 421 | 83.2 | 63 | 16.8 | 4.6 |
| Gender | | | | | |
| Male | 166 | 77.7 | 28 | 22.3 | 8.1 |
| Female | 255 | 88.2 | 35 | 11.8 | 4.4 |
| Age | | | | | |
| 18-34 | 155 | 79.6 | 26 | 20.4 | 8.2 |
| 35-54 | 202 | 85.8 | 29 | 14.2 | 5.6 |
| 55-64 | 64 | 86.1 | 8 | 13.9 | 11.6 |
| Education | | | | | |
| ≤High School | 78 | 72.3 | 21 | 27.7 | 12.1 |
| >High School | 343 | 87.1 | 41 | 12.9 | 4.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Health Insurance Coverage, Aged 18 and Older

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|----|------|-------------------|
| Total | 547 | 85.4 | 66 | 14.6 | 4.0 |
| Gender | | | | | |
| Male | 216 | 80.7 | 28 | 19.3 | 7.1 |
| Female | 331 | 89.5 | 38 | 10.5 | 3.8 |
| Age | | | | | |
| 18-34 | 155 | 79.6 | 26 | 20.4 | 8.2 |
| 35-54 | 202 | 85.8 | 29 | 14.2 | 5.6 |
| ≥55 | 176 | 92.0 | 11 | 8.0 | 5.8 |
| Education | | | | | |
| ≤High School | 123 | 78.0 | 22 | 22.0 | 9.7 |
| >High School | 421 | 88.5 | 43 | 11.5 | 3.8 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Reported Having a Routine Physical Exam in Past Two Years

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|----|------|-------------------|
| Total | 539 | 88.0 | 70 | 12.0 | 3.1 |
| Gender | | | | | |
| Male | 206 | 84.5 | 35 | 15.5 | 5.5 |
| Female | 333 | 91.0 | 35 | 9.0 | 3.3 |
| Age | | | | | |
| 18-34 | 151 | 83.0 | 28 | 17.0 | 6.7 |
| 35-54 | 201 | 88.1 | 29 | 11.9 | 4.7 |
| ≥55 | 176 | 94.5 | 12 | 5.5 | 3.3 |
| Education | | | | | |
| ≤High School | 136 | 90.5 | 12 | 9.5 | 6.4 |
| >High School | 399 | 86.8 | 58 | 13.2 | 3.6 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Time in Past 12 Months when Needed Medical Care but Could Not Get It

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 49 | 10.3 | 567 | 89.7 | 3.4 |
| Gender | | | | | |
| Male | 17 | 10.5 | 229 | 89.5 | 5.7 |
| Female | 32 | 10.2 | 338 | 89.8 | 3.9 |
| Age | | | | | |
| 18-34 | 24 | 18.7 | 157 | 81.3 | 7.8 |
| 35-54 | 18 | 7.8 | 215 | 92.2 | 4.0 |
| ≥55 | 7 | 3.0 | 181 | 97.0 | 2.6 |
| Education | | | | | |
| ≤High School | 21 | 19.5 | 127 | 80.5 | 9.1 |
| >High School | 27 | 6.4 | 437 | 93.6 | 2.7 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 493 | 78.3 | 120 | 21.7 | 4.0 |
| Gender | | | | | |
| Male | 208 | 81.3 | 36 | 18.7 | 6.1 |
| Female | 285 | 75.6 | 84 | 24.4 | 5.3 |
| Age | | | | | |
| 18-34 | 160 | 86.9 | 20 | 13.1 | 6.5 |
| 35-54 | 181 | 74.0 | 51 | 26.0 | 6.8 |
| ≥55 | 143 | 73.8 | 44 | 26.2 | 7.7 |
| Education | | | | | |
| ≤High School | 83 | 54.3 | 64 | 45.7 | 9.7 |
| >High School | 407 | 88.2 | 55 | 11.8 | 3.4 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Physician Diagnosed Diabetes

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 34 | 5.9 | 582 | 94.1 | 2.2 |
| Gender | | | | | |
| Male | 18 | 7.5 | 228 | 92.5 | 3.7 |
| Female | 16 | 4.4 | 354 | 95.6 | 2.6 |
| Age | | | | | |
| 18-34 | 1 | 0.7 | 180 | 99.3 | 1.3 |
| 35-54 | 11 | 5.7 | 222 | 94.3 | 3.6 |
| ≥55 | 22 | 13.4 | 166 | 86.6 | 6.3 |
| Education | | | | | |
| ≤High School | 17 | 12.1 | 131 | 87.9 | 6.3 |
| >High School | 17 | 3.3 | 447 | 96.7 | 1.7 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Lifetime Asthma Diagnosed by a Medical Professional

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 77 | 12.7 | 539 | 87.3 | 3.1 |
| Gender | | | | | |
| Male | 30 | 13.1 | 216 | 86.9 | 5.1 |
| Female | 47 | 12.4 | 323 | 87.6 | 3.8 |
| Age | | | | | |
| 18-34 | 23 | 14.3 | 158 | 85.7 | 5.9 |
| 35-54 | 24 | 8.9 | 209 | 91.1 | 3.8 |
| ≥55 | 28 | 15.8 | 160 | 84.2 | 6.7 |
| Education | | | | | |
| ≤High School | 20 | 12.7 | 128 | 87.3 | 6.5 |
| >High School | 57 | 12.8 | 407 | 87.2 | 3.5 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Current Asthma Diagnosed by a Medical Professional

New York

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 50 | 8.0 | 564 | 92.0 | 2.6 |
| Gender | | | | | |
| Male | 20 | 8.7 | 225 | 91.3 | 4.4 |
| Female | 30 | 7.3 | 339 | 92.7 | 2.9 |
| Age | | | | | |
| 18-34 | 12 | 7.8 | 169 | 92.2 | 4.5 |
| 35-54 | 16 | 5.8 | 215 | 94.2 | 3.0 |
| ≥55 | 20 | 11.0 | 168 | 89.0 | 6.0 |
| Education | | | | | |
| ≤High School | 16 | 10.5 | 132 | 89.5 | 6.2 |
| >High School | 34 | 7.0 | 428 | 93.0 | 2.6 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Arthritis Diagnosed by a Medical Professional

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 145 | 21.9 | 467 | 78.1 | 3.7 |
| Gender | | | | | |
| Male | 47 | 17.3 | 197 | 82.7 | 5.2 |
| Female | 98 | 25.9 | 270 | 74.1 | 5.2 |
| Age | | | | | |
| 18-34 | 10 | 5.2 | 171 | 94.8 | 3.7 |
| 35-54 | 41 | 16.6 | 190 | 83.4 | 5.2 |
| ≥55 | 92 | 51.8 | 95 | 48.2 | 8.5 |
| Education | | | | | |
| ≤High School | 51 | 29.9 | 97 | 70.1 | 8.5 |
| >High School | 92 | 18.4 | 368 | 81.6 | 3.9 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)*New York*

| Demographic Groups | n¹ | Doctor Diagnosed %² | C.I.³ | n | Possible Arthritis (CJS Only) % | C.I. | n | No Arthritis or CJS % | C.I. |
|---------------------------|----------------------|---------------------------------------|-------------------------|----------|--|-------------|----------|------------------------------|-------------|
| Total | 145 | 22.0 | 3.8 | 80 | 13.4 | 3.3 | 384 | 64.6 | 4.5 |
| Gender | | | | | | | | | |
| Male | 47 | 17.4 | 5.2 | 35 | 14.8 | 5.5 | 161 | 67.8 | 6.8 |
| Female | 98 | 26.0 | 5.3 | 45 | 12.1 | 3.9 | 223 | 61.9 | 5.8 |
| Age | | | | | | | | | |
| 18-34 | 10 | 5.2 | 3.7 | 24 | 15.6 | 6.6 | 147 | 79.2 | 7.2 |
| 35-54 | 41 | 16.6 | 5.2 | 36 | 14.8 | 5.3 | 154 | 68.6 | 6.8 |
| ≥55 | 92 | 52.6 | 8.6 | 17 | 8.1 | 4.1 | 76 | 39.3 | 8.4 |
| Education | | | | | | | | | |
| ≤High School | 51 | 30.2 | 8.6 | 20 | 12.9 | 6.9 | 76 | 57.0 | 9.6 |
| >High School | 92 | 18.5 | 4.0 | 60 | 13.7 | 3.7 | 306 | 67.8 | 4.9 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution²Weighted Percent³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Smoked 100 Cigarettes in Lifetime

New York

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 285 | 44.7 | 329 | 55.3 | 4.6 |
| Gender | | | | | |
| Male | 118 | 48.2 | 128 | 51.8 | 7.3 |
| Female | 167 | 41.5 | 201 | 58.5 | 5.7 |
| Age | | | | | |
| 18-34 | 65 | 36.1 | 115 | 63.9 | 8.0 |
| 35-54 | 110 | 44.7 | 123 | 55.3 | 7.3 |
| ≥55 | 101 | 55.6 | 86 | 44.4 | 8.5 |
| Education | | | | | |
| ≤High School | 63 | 39.1 | 85 | 60.9 | 9.3 |
| >High School | 219 | 46.7 | 243 | 53.3 | 5.2 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Current Smoker

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 114 | 19.3 | 500 | 80.7 | 3.7 |
| Gender | | | | | |
| Male | 40 | 18.7 | 206 | 81.3 | 5.9 |
| Female | 74 | 19.9 | 294 | 80.1 | 4.7 |
| Age | | | | | |
| 18-34 | 37 | 21.1 | 143 | 78.9 | 6.8 |
| 35-54 | 49 | 19.4 | 184 | 80.6 | 5.6 |
| ≥55 | 25 | 16.5 | 162 | 83.5 | 7.2 |
| Education | | | | | |
| ≤High School | 33 | 22.4 | 115 | 77.6 | 8.1 |
| >High School | 80 | 18.0 | 382 | 82.0 | 4.1 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Status of Cigarette Smoking*New York*

| Demographic Groups | n¹ | Everyday Smoker %² | C.I.³ | n | Someday Smoker % | C.I. | n | Former or Never Smoked % | C.I. |
|---------------------------|----------------------|--|-------------------------|----------|-----------------------------|-------------|----------|---|-------------|
| Total | 72 | 12.3 | 3.1 | 42 | 7.0 | 2.5 | 500 | 80.7 | 3.7 |
| Gender | | | | | | | | | |
| Male | 25 | 11.0 | 4.6 | 15 | 7.7 | 4.3 | 206 | 81.3 | 5.9 |
| Female | 47 | 13.6 | 4.1 | 27 | 6.3 | 2.7 | 294 | 80.1 | 4.7 |
| Age | | | | | | | | | |
| 18-34 | 21 | 13.4 | 5.9 | 16 | 7.6 | 4.1 | 143 | 78.9 | 6.8 |
| 35-54 | 32 | 13.1 | 4.8 | 17 | 6.3 | 3.4 | 184 | 80.6 | 5.6 |
| ≥55 | 17 | 9.3 | 4.9 | 8 | 7.2 | 5.9 | 162 | 83.5 | 7.2 |
| Education | | | | | | | | | |
| ≤High School | 22 | 14.4 | 6.5 | 11 | 8.0 | 5.6 | 115 | 77.6 | 8.1 |
| >High School | 49 | 11.4 | 3.4 | 31 | 6.6 | 2.7 | 382 | 82.0 | 4.1 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution²Weighted Percent³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Home Smoking Policies*New York*

| Demographic Groups | n¹ | Not Allowed %² | C.I.³ | n | Allowed Some Places % | C.I. | n | Allowed Anywhere % | C.I. |
|---------------------------|----------------------|----------------------------------|-------------------------|----------|------------------------------|-------------|----------|---------------------------|-------------|
| Total | 377 | 61.4 | 4.6 | 79 | 12.8 | 3.0 | 157 | 25.9 | 4.2 |
| Gender | | | | | | | | | |
| Male | 139 | 57.3 | 7.3 | 29 | 11.7 | 4.4 | 77 | 31.0 | 6.9 |
| Female | 238 | 64.9 | 5.6 | 50 | 13.8 | 4.1 | 80 | 21.3 | 4.8 |
| Age | | | | | | | | | |
| 18-34 | 113 | 60.7 | 8.4 | 30 | 15.8 | 5.8 | 38 | 23.6 | 7.7 |
| 35-54 | 136 | 60.2 | 7.2 | 33 | 14.3 | 5.2 | 62 | 25.5 | 6.3 |
| ≥55 | 117 | 62.5 | 8.3 | 16 | 7.5 | 3.9 | 54 | 30.0 | 8.0 |
| Education | | | | | | | | | |
| ≤High School | 99 | 62.8 | 9.8 | 14 | 8.8 | 4.9 | 34 | 28.4 | 9.6 |
| >High School | 275 | 60.6 | 5.1 | 65 | 14.5 | 3.7 | 122 | 24.8 | 4.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution²Weighted Percent³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Home Smoking

New York

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 147 | 26.0 | 469 | 74.0 | 4.3 |
| Gender | | | | | |
| Male | 59 | 27.2 | 187 | 72.8 | 7.0 |
| Female | 88 | 25.0 | 282 | 75.0 | 5.2 |
| Age | | | | | |
| 18-34 | 52 | 32.9 | 129 | 67.1 | 8.4 |
| 35-54 | 57 | 21.8 | 176 | 78.2 | 5.9 |
| ≥55 | 37 | 23.6 | 151 | 76.4 | 7.7 |
| Education | | | | | |
| ≤High School | 36 | 30.3 | 112 | 69.7 | 9.8 |
| >High School | 110 | 24.3 | 354 | 75.7 | 4.5 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)

New York

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 97 | 16.0 | 508 | 84.0 | 3.3 |
| Gender | | | | | |
| Male | 55 | 21.4 | 186 | 78.6 | 5.7 |
| Female | 42 | 11.4 | 322 | 88.6 | 3.8 |
| Age | | | | | |
| 18-34 | 52 | 27.7 | 124 | 72.3 | 7.3 |
| 35-54 | 35 | 12.8 | 194 | 87.2 | 4.6 |
| ≥55 | 10 | 6.1 | 176 | 93.9 | 4.3 |
| Education | | | | | |
| ≤High School | 16 | 12.7 | 129 | 87.3 | 6.6 |
| >High School | 81 | 17.4 | 376 | 82.6 | 3.9 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)

New York

| Demographic Groups | n ¹ | % ² | n | No % | C.I. ³ |
|--------------------|----------------|----------------|-----|------|-------------------|
| Total | 51 | 7.6 | 557 | 92.4 | 2.3 |
| Gender | | | | | |
| Male | 19 | 7.2 | 222 | 92.8 | 3.4 |
| Female | 32 | 8.0 | 335 | 92.0 | 3.1 |
| Age | | | | | |
| 18-34 | 14 | 8.3 | 163 | 91.7 | 4.5 |
| 35-54 | 19 | 6.4 | 212 | 93.6 | 3.1 |
| ≥55 | 16 | 8.5 | 170 | 91.5 | 4.5 |
| Education | | | | | |
| ≤High School | 8 | 5.0 | 137 | 95.0 | 3.7 |
| >High School | 43 | 8.7 | 417 | 91.3 | 2.8 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Overweight or Obese, BMI(Body Mass Index) 25 or More

New York

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 237 | 42.3 | 347 | 57.7 | 4.7 |
| Gender | | | | | |
| Male | 127 | 51.3 | 115 | 48.7 | 7.4 |
| Female | 110 | 34.0 | 232 | 66.0 | 5.9 |
| Age | | | | | |
| 18-34 | 48 | 31.0 | 126 | 69.0 | 8.1 |
| 35-54 | 102 | 48.7 | 120 | 51.3 | 7.6 |
| ≥55 | 84 | 49.0 | 95 | 51.0 | 8.8 |
| Education | | | | | |
| ≤High School | 72 | 52.0 | 64 | 48.0 | 10.2 |
| >High School | 164 | 38.3 | 283 | 61.7 | 5.2 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+)

New York

| Demographic Groups | n¹ | Underweight\ Normal %² | C.I.³ | n | Overweight % | C.I. | n | Obese % | C.I. |
|---------------------------|----------------------|--|-------------------------|----------|---------------------|-------------|----------|----------------|-------------|
| Total | 347 | 57.7 | 4.7 | 163 | 28.6 | 4.3 | 74 | 13.7 | 3.4 |
| Gender | | | | | | | | | |
| Male | 115 | 48.7 | 7.4 | 100 | 39.6 | 7.1 | 27 | 11.7 | 5.0 |
| Female | 232 | 66.0 | 5.9 | 63 | 18.5 | 4.8 | 47 | 15.5 | 4.7 |
| Age | | | | | | | | | |
| 18-34 | 126 | 69.0 | 8.1 | 36 | 20.0 | 6.4 | 12 | 10.9 | 6.5 |
| 35-54 | 120 | 51.3 | 7.6 | 71 | 35.6 | 7.4 | 31 | 13.1 | 4.8 |
| ≥55 | 95 | 51.0 | 8.8 | 53 | 30.3 | 8.3 | 31 | 18.7 | 6.8 |
| Education | | | | | | | | | |
| ≤High School | 64 | 48.0 | 10.2 | 38 | 24.6 | 8.1 | 34 | 27.4 | 9.2 |
| >High School | 283 | 61.7 | 5.2 | 125 | 30.3 | 5.0 | 39 | 8.0 | 2.7 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had Mammography Screening, Women Aged 40 Years and Older

New York

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Female | 192 | 93.6 | 19 | 6.4 | 3.0 |
| Age | | | | | |
| 40-64 | 131 | 93.4 | 14 | 6.6 | 3.7 |
| ≥65 | 61 | 94.2 | 5 | 5.8 | 5.1 |
| Education | | | | | |
| ≤High School | 58 | 95.6 | 4 | 4.4 | 4.7 |
| >High School | 133 | 92.4 | 15 | 7.6 | 4.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older

New York

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Female | 160 | 79.4 | 49 | 20.6 | 5.9 |
| Age | | | | | |
| 40-64 | 112 | 80.1 | 32 | 19.9 | 7.1 |
| ≥65 | 48 | 77.6 | 17 | 22.4 | 10.9 |
| Education | | | | | |
| ≤High School | 55 | 91.1 | 6 | 8.9 | 7.9 |
| >High School | 105 | 73.3 | 42 | 26.7 | 7.9 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health
Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger
New York

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 116 | 25.8 | 359 | 74.2 | 4.8 |
| Gender | | | | | |
| Male | 32 | 20.9 | 160 | 79.1 | 7.6 |
| Female | 84 | 30.4 | 199 | 69.6 | 6.2 |
| Age | | | | | |
| 18-34 | 62 | 36.7 | 115 | 63.3 | 8.6 |
| 35-54 | 48 | 20.2 | 177 | 79.8 | 5.8 |
| 55-64 | 6 | 9.5 | 67 | 90.5 | 8.6 |
| Education | | | | | |
| ≤High School | 41 | 43.3 | 57 | 56.7 | 12.0 |
| >High School | 75 | 19.8 | 301 | 80.2 | 4.6 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Sexually Active, Age 18-64 Years Old

New York

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 328 | 80.0 | 107 | 20.0 | 4.4 |
| Gender | | | | | |
| Male | 145 | 80.8 | 40 | 19.2 | 6.8 |
| Female | 183 | 79.3 | 67 | 20.7 | 5.4 |
| Age | | | | | |
| 18-34 | 138 | 84.8 | 28 | 15.2 | 6.5 |
| 35-54 | 154 | 80.2 | 50 | 19.8 | 6.2 |
| 55-64 | 36 | 64.8 | 29 | 35.2 | 13.8 |
| Education | | | | | |
| ≤High School | 68 | 79.5 | 24 | 20.5 | 10.1 |
| >High School | 259 | 80.2 | 83 | 19.8 | 4.7 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger

New York

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 141 | 38.0 | 213 | 62.0 | 5.8 |
| Gender | | | | | |
| Male | 66 | 40.6 | 89 | 59.4 | 8.8 |
| Female | 75 | 35.3 | 124 | 64.7 | 7.5 |
| Age | | | | | |
| 18-34 | 76 | 50.1 | 73 | 49.9 | 9.3 |
| 35-54 | 61 | 32.1 | 103 | 67.9 | 7.8 |
| 55-64 | 4 | 10.3 | 37 | 89.7 | 11.0 |
| Education | | | | | |
| ≤High School | 31 | 43.3 | 41 | 56.7 | 13.2 |
| >High School | 109 | 35.9 | 172 | 64.1 | 6.2 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)

New York

| Demographic Groups | Use BC | | | n | No BC at risk | | | No BC | | |
|--------------------|----------------|----------------|-------------------|----|---------------|------|-----|-------|------|--|
| | n ¹ | % ² | C.I. ³ | | % | C.I. | n | % | C.I. | |
| Total | 193 | 58.3 | 6.0 | 40 | 13.0 | 4.1 | 110 | 28.7 | 5.4 | |
| Gender | | | | | | | | | | |
| Male | 87 | 53.6 | 8.5 | 22 | 13.0 | 5.7 | 67 | 33.4 | 7.8 | |
| Female | 106 | 64.8 | 8.2 | 18 | 13.0 | 5.9 | 43 | 22.2 | 7.1 | |
| Age | | | | | | | | | | |
| 18-34 | 116 | 68.1 | 7.8 | 16 | 10.0 | 5.0 | 42 | 21.9 | 6.8 | |
| 35-59 | 77 | 46.1 | 8.6 | 24 | 16.7 | 6.7 | 68 | 37.2 | 8.5 | |
| Education | | | | | | | | | | |
| ≤High School | 41 | 66.2 | 13.1 | 9 | 16.1 | 10.3 | 15 | 17.7 | 9.8 | |
| >High School | 151 | 55.7 | 6.6 | 31 | 12.0 | 4.3 | 95 | 32.3 | 6.3 | |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Physician Diagnosed Heart Attack, Angina or Stroke

New York

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 38 | 5.6 | 565 | 94.4 | 2.1 |
| Gender | | | | | |
| Male | 22 | 7.3 | 220 | 92.7 | 3.7 |
| Female | 16 | 4.1 | 345 | 95.9 | 2.3 |
| Age | | | | | |
| 18-64 | 11 | 2.5 | 466 | 97.5 | 1.8 |
| ≥65 | 27 | 23.5 | 87 | 76.5 | 9.0 |
| Education | | | | | |
| ≤High School | 14 | 9.2 | 126 | 90.8 | 5.6 |
| >High School | 24 | 4.2 | 435 | 95.8 | 2.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina)

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 32 | 4.6 | 570 | 95.4 | 1.9 |
| Gender | | | | | |
| Male | 19 | 6.6 | 222 | 93.4 | 3.5 |
| Female | 13 | 2.9 | 348 | 97.1 | 1.6 |
| Age | | | | | |
| 18-64 | 10 | 2.4 | 466 | 97.6 | 1.8 |
| ≥65 | 22 | 17.4 | 92 | 82.6 | 7.3 |
| Education | | | | | |
| ≤High School | 13 | 7.7 | 127 | 92.3 | 4.8 |
| >High School | 19 | 3.4 | 439 | 96.6 | 1.9 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older

New York

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Male | 89 | 68.0 | 42 | 32.0 | 9.3 |
| Age | | | | | |
| 40-64 | 51 | 61.1 | 36 | 38.9 | 11.7 |
| ≥65 | 38 | 88.0 | 6 | 12.0 | 10.5 |
| Education | | | | | |
| ≤High School | 23 | 66.8 | 9 | 33.2 | 20.7 |
| >High School | 65 | 68.0 | 33 | 32.0 | 10.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|----|------|-------------------|
| Male | 76 | 59.7 | 53 | 40.3 | 9.8 |
| Age | | | | | |
| 40-64 | 43 | 53.5 | 43 | 46.5 | 12.1 |
| ≥65 | 33 | 78.1 | 10 | 21.9 | 14.3 |
| Education | | | | | |
| ≤High School | 19 | 58.4 | 12 | 41.6 | 21.2 |
| >High School | 56 | 59.7 | 41 | 40.3 | 11.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had a Digital Rectal Exam, Men Aged 40 and Older

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|----|------|-------------------|
| Male | 121 | 84.9 | 19 | 15.1 | 7.5 |
| Age | | | | | |
| 40-64 | 77 | 81.0 | 16 | 19.0 | 9.7 |
| ≥65 | 44 | 96.3 | 3 | 3.7 | 4.3 |
| Education | | | | | |
| ≤High School | 27 | 78.5 | 6 | 21.5 | 19.3 |
| >High School | 93 | 86.9 | 13 | 13.1 | 7.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Used a Blood Stool Test at Home, Aged 50 and Older

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 82 | 35.2 | 156 | 64.8 | 7.2 |
| Gender | | | | | |
| Male | 41 | 42.0 | 52 | 58.0 | 11.8 |
| Female | 41 | 29.8 | 104 | 70.2 | 8.7 |
| Age | | | | | |
| 50-64 | 39 | 31.3 | 85 | 68.7 | 9.6 |
| ≥65 | 43 | 40.4 | 71 | 59.6 | 10.5 |
| Education | | | | | |
| ≤High School | 21 | 33.3 | 45 | 66.7 | 13.6 |
| >High School | 61 | 36.8 | 109 | 63.2 | 8.4 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 38 | 17.3 | 199 | 82.7 | 5.8 |
| Gender | | | | | |
| Male | 19 | 21.4 | 73 | 78.6 | 10.2 |
| Female | 19 | 14.1 | 126 | 85.9 | 6.5 |
| Age | | | | | |
| 50-64 | 16 | 14.2 | 107 | 85.8 | 7.6 |
| ≥65 | 22 | 21.4 | 92 | 78.6 | 8.9 |
| Education | | | | | |
| ≤High School | 10 | 17.3 | 56 | 82.7 | 11.1 |
| >High School | 28 | 17.6 | 141 | 82.4 | 6.8 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|-----|------|-------------------|
| Total | 56 | 26.1 | 181 | 73.9 | 6.8 |
| Gender | | | | | |
| Male | 29 | 31.0 | 63 | 69.0 | 11.3 |
| Female | 27 | 22.2 | 118 | 77.8 | 8.4 |
| Age | | | | | |
| 50-64 | 23 | 21.0 | 100 | 79.0 | 8.9 |
| ≥65 | 33 | 32.8 | 81 | 67.2 | 10.4 |
| Education | | | | | |
| ≤High School | 15 | 26.9 | 51 | 73.1 | 13.2 |
| >High School | 41 | 26.1 | 128 | 73.9 | 7.9 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older

New York

| Demographic Groups | n ¹ | Yes % ² | n | No % | C.I. ³ |
|--------------------|----------------|--------------------|----|------|-------------------|
| Total | 139 | 58.6 | 98 | 41.4 | 7.4 |
| Gender | | | | | |
| Male | 56 | 57.9 | 36 | 42.1 | 12.1 |
| Female | 83 | 59.3 | 62 | 40.7 | 9.0 |
| Age | | | | | |
| 50-64 | 64 | 53.0 | 60 | 47.0 | 10.4 |
| ≥65 | 75 | 66.1 | 38 | 33.9 | 10.0 |
| Education | | | | | |
| ≤High School | 36 | 53.3 | 29 | 46.7 | 14.4 |
| >High School | 103 | 62.3 | 67 | 37.7 | 8.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older

New York

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 135 | 57.6 | 101 | 42.4 | 7.4 |
| Gender | | | | | |
| Male | 52 | 55.5 | 39 | 44.5 | 12.2 |
| Female | 83 | 59.3 | 62 | 40.7 | 9.0 |
| Age | | | | | |
| 50-64 | 63 | 52.4 | 61 | 47.6 | 10.4 |
| ≥65 | 72 | 64.5 | 40 | 35.5 | 10.1 |
| Education | | | | | |
| ≤High School | 35 | 52.4 | 30 | 47.6 | 14.4 |
| >High School | 100 | 61.2 | 69 | 38.8 | 8.3 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older

New York

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---------------------------|----------------------|--------------------------|----------|-------------|-------------------------|
| Total | 145 | 61.6 | 94 | 38.4 | 7.3 |
| Gender | | | | | |
| Male | 57 | 58.7 | 36 | 41.3 | 12.0 |
| Female | 88 | 63.9 | 58 | 36.1 | 8.7 |
| Age | | | | | |
| 50-64 | 66 | 54.3 | 59 | 45.7 | 10.3 |
| ≥65 | 79 | 71.2 | 35 | 28.8 | 9.4 |
| Education | | | | | |
| ≤High School | 40 | 59.9 | 26 | 40.1 | 14.2 |
| >High School | 105 | 63.3 | 66 | 36.7 | 8.2 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval