# Expanded BRFSS, 2003

## Albany Locality Tables

TABLE OF CONTENTS:

**New York State Department Of Health**

Expanded BRFSS, 2003 - New York State Department of Health

# Self-rated Health Status

*Albany*

| Demographic Groups | n[1] | Fair or Poor %[2] | n | Good to Excellent % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 86 | 14.5 | 551 | 85.5 | 3.4 |
| Gender | | | | | |
| Male | 33 | 13.8 | 219 | 86.2 | 5.3 |
| Female | 53 | 15.1 | 332 | 84.9 | 4.3 |
| Age | | | | | |
| 18-34 | 11 | 11.2 | 152 | 88.8 | 7.1 |
| 35-54 | 30 | 12.1 | 214 | 87.9 | 4.9 |
| ≥55 | 44 | 21.1 | 170 | 78.9 | 6.0 |
| Education | | | | | |
| ≤High School | 44 | 24.9 | 140 | 75.1 | 7.3 |
| >High School | 42 | 10.2 | 410 | 89.8 | 3.7 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health
## Reported Number of Physical Distress Days in the Past Month
### *Albany*

| Demographic Groups | n[1] | ≥14 days %[2] | n | < 14 days % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 54 | 8.6 | 580 | 91.4 | 2.8 |
| Gender | | | | | |
| Male | 18 | 8.2 | 232 | 91.8 | 4.5 |
| Female | 36 | 9.1 | 348 | 90.9 | 3.4 |
| Age | | | | | |
| 18-34 | 5 | 4.5 | 161 | 95.5 | 5.1 |
| 35-54 | 25 | 10.0 | 218 | 90.0 | 4.5 |
| ≥55 | 23 | 11.7 | 187 | 88.3 | 5.0 |
| Education | | | | | |
| ≤High School | 18 | 9.5 | 163 | 90.5 | 5.1 |
| >High School | 36 | 8.3 | 416 | 91.7 | 3.3 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health
## Reported Number of Mental Distress Days in the Past Month
### *Albany*

| Demographic Groups | n[1] | ≥14 days %[2] | n | < 14 days % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 61 | 10.6 | 576 | 89.4 | 3.0 |
| Gender | | | | | |
| Male | 14 | 6.7 | 236 | 93.3 | 4.3 |
| Female | 47 | 14.0 | 340 | 86.0 | 4.2 |
| Age | | | | | |
| 18-34 | 19 | 14.9 | 146 | 85.1 | 7.0 |
| 35-54 | 28 | 11.2 | 212 | 88.8 | 4.7 |
| ≥55 | 14 | 6.2 | 202 | 93.8 | 3.6 |
| Education | | | | | |
| ≤High School | 20 | 13.0 | 164 | 87.0 | 6.2 |
| >High School | 41 | 9.6 | 411 | 90.4 | 3.4 |

[1] Percents based on row denominators of less than 50 are unstable and should be used with caution
[2] Weighted Percent
[3] 95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

# Health Insurance Coverage, 18-64 Years of Age

## *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 469 | 90.0 | 42 | 10.0 | 3.4 |
| Gender | | | | | |
| Male | 185 | 86.2 | 24 | 13.8 | 6.1 |
| Female | 284 | 93.7 | 18 | 6.3 | 3.1 |
| Age | | | | | |
| 18-34 | 140 | 81.9 | 26 | 18.1 | 7.4 |
| 35-54 | 232 | 95.7 | 11 | 4.3 | 2.7 |
| 55-64 | 97 | 94.5 | 5 | 5.5 | 4.9 |
| Education | | | | | |
| ≤High School | 114 | 87.6 | 17 | 12.4 | 6.2 |
| >High School | 355 | 90.8 | 25 | 9.2 | 4.1 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

# Health Insurance Coverage, Aged 18 and Older
## *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 597 | 91.7 | 44 | 8.3 | 2.8 |
| Gender | | | | | |
| Male | 227 | 88.3 | 24 | 11.7 | 5.2 |
| Female | 370 | 94.7 | 20 | 5.3 | 2.5 |
| Age | | | | | |
| 18-34 | 140 | 81.9 | 26 | 18.1 | 7.4 |
| 35-54 | 232 | 95.7 | 11 | 4.3 | 2.7 |
| ≥55 | 210 | 96.7 | 7 | 3.3 | 2.5 |
| Education | | | | | |
| ≤High School | 167 | 90.3 | 19 | 9.7 | 4.6 |
| >High School | 429 | 92.3 | 25 | 7.7 | 3.5 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Reported Having a Routine Physical Exam in Past Two Years
### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 579 | 88.8 | 61 | 11.2 | 2.9 |
| Gender | | | | | |
| Male | 215 | 83.4 | 36 | 16.6 | 5.3 |
| Female | 364 | 93.5 | 25 | 6.5 | 2.8 |
| Age | | | | | |
| 18-34 | 144 | 85.8 | 21 | 14.2 | 6.3 |
| 35-54 | 215 | 85.4 | 29 | 14.6 | 5.3 |
| ≥55 | 204 | 94.7 | 11 | 5.3 | 3.2 |
| Education | | | | | |
| ≤High School | 175 | 94.3 | 9 | 5.7 | 3.8 |
| >High School | 403 | 86.5 | 52 | 13.5 | 3.8 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Time in Past 12 Months when Needed Medical Care but Could Not Get It
### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 32 | 5.7 | 611 | 94.3 | 2.3 |
| Gender | | | | | |
| Male | 9 | 4.1 | 244 | 95.9 | 3.1 |
| Female | 23 | 7.1 | 367 | 92.9 | 3.3 |
| Age | | | | | |
| 18-34 | 16 | 10.2 | 150 | 89.8 | 5.4 |
| 35-54 | 12 | 6.0 | 232 | 94.0 | 4.1 |
| ≥55 | 3 | 1.1 | 214 | 98.9 | 1.4 |
| Education | | | | | |
| ≤High School | 9 | 7.3 | 177 | 92.7 | 5.6 |
| >High School | 23 | 5.0 | 433 | 95.0 | 2.2 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days
### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 518 | 80.6 | 123 | 19.4 | 3.6 |
| Gender | | | | | |
| Male | 212 | 82.8 | 41 | 17.2 | 5.7 |
| Female | 306 | 78.6 | 82 | 21.4 | 4.6 |
| Age | | | | | |
| 18-34 | 138 | 81.4 | 28 | 18.6 | 7.5 |
| 35-54 | 201 | 82.0 | 43 | 18.0 | 5.6 |
| ≥55 | 167 | 78.2 | 49 | 21.8 | 5.9 |
| Education | | | | | |
| ≤High School | 130 | 68.3 | 56 | 31.7 | 7.9 |
| >High School | 388 | 85.8 | 66 | 14.2 | 3.8 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Physician Diagnosed Diabetes

### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 40 | 6.3 | 603 | 93.7 | 2.2 |
| Gender | | | | | |
| Male | 16 | 5.4 | 237 | 94.6 | 2.8 |
| Female | 24 | 7.1 | 366 | 92.9 | 3.2 |
| Age | | | | | |
| 18-34 | 2 | 1.2 | 164 | 98.8 | 1.8 |
| 35-54 | 14 | 6.5 | 230 | 93.5 | 4.0 |
| ≥55 | 24 | 11.5 | 193 | 88.5 | 4.7 |
| Education | | | | | |
| ≤High School | 14 | 9.2 | 172 | 90.8 | 5.3 |
| >High School | 26 | 5.1 | 430 | 94.9 | 2.1 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Lifetime Asthma Diagnosed by a Medical Professional
### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 91 | 14.6 | 552 | 85.4 | 3.4 |
| Gender | | | | | |
| Male | 23 | 10.8 | 230 | 89.2 | 5.1 |
| Female | 68 | 18.0 | 322 | 82.0 | 4.5 |
| Age | | | | | |
| 18-34 | 28 | 19.2 | 138 | 80.8 | 7.8 |
| 35-54 | 33 | 12.6 | 211 | 87.4 | 4.5 |
| ≥55 | 29 | 12.9 | 188 | 87.1 | 4.8 |
| Education | | | | | |
| ≤High School | 22 | 10.8 | 164 | 89.2 | 4.9 |
| >High School | 69 | 16.2 | 387 | 83.8 | 4.3 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Current Asthma Diagnosed by a Medical Professional
### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 60 | 9.8 | 580 | 90.2 | 2.9 |
| Gender | | | | | |
| Male | 12 | 5.9 | 241 | 94.1 | 4.0 |
| Female | 48 | 13.2 | 339 | 86.8 | 4.1 |
| Age | | | | | |
| 18-34 | 18 | 13.3 | 147 | 86.7 | 7.0 |
| 35-54 | 23 | 9.0 | 221 | 91.0 | 3.9 |
| ≥55 | 19 | 7.7 | 196 | 92.3 | 3.6 |
| Education | | | | | |
| ≤High School | 14 | 6.2 | 170 | 93.8 | 3.6 |
| >High School | 46 | 11.2 | 409 | 88.8 | 3.8 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

# Arthritis Diagnosed by a Medical Professional
## *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 184 | 26.5 | 457 | 73.5 | 3.7 |
| Gender | | | | | |
| Male | 58 | 21.4 | 195 | 78.6 | 5.3 |
| Female | 126 | 31.0 | 262 | 69.0 | 5.2 |
| Age | | | | | |
| 18-34 | 9 | 4.6 | 156 | 95.4 | 3.1 |
| 35-54 | 59 | 22.3 | 185 | 77.7 | 5.9 |
| ≥55 | 110 | 52.0 | 107 | 48.0 | 7.2 |
| Education | | | | | |
| ≤High School | 70 | 35.4 | 116 | 64.6 | 7.7 |
| >High School | 114 | 22.8 | 340 | 77.2 | 4.1 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

*Albany*

| Demographic Groups | n[1] | Doctor Diagnosed %[2] | C.I.[3] | n | Possible Arthritis (CJS Only) % | C.I. | n | No Arthritis or CJS % | C.I. |
|---|---|---|---|---|---|---|---|---|---|
| Total | 184 | 26.5 | 3.7 | 78 | 14.2 | 3.5 | 379 | 59.3 | 4.4 |
| Gender | | | | | | | | | |
| Male | 58 | 21.4 | 5.3 | 35 | 15.5 | 5.5 | 160 | 63.1 | 6.6 |
| Female | 126 | 31.0 | 5.2 | 43 | 12.9 | 4.4 | 219 | 56.0 | 5.7 |
| Age | | | | | | | | | |
| 18-34 | 9 | 4.6 | 3.1 | 19 | 16.5 | 8.0 | 137 | 78.9 | 8.2 |
| 35-54 | 59 | 22.3 | 5.9 | 40 | 17.2 | 5.3 | 145 | 60.5 | 6.9 |
| ≥55 | 110 | 52.0 | 7.2 | 18 | 8.9 | 4.2 | 89 | 39.1 | 7.0 |
| Education | | | | | | | | | |
| ≤High School | 70 | 35.4 | 7.7 | 23 | 12.6 | 5.3 | 93 | 52.0 | 8.1 |
| >High School | 114 | 22.8 | 4.1 | 55 | 14.8 | 4.4 | 285 | 62.4 | 5.2 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

# Ever Smoked 100 Cigarettes in Lifetime
## *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 303 | 45.3 | 339 | 54.7 | 4.4 |
| Gender | | | | | |
| Male | 119 | 46.1 | 134 | 53.9 | 6.8 |
| Female | 184 | 44.5 | 205 | 55.5 | 5.6 |
| Age | | | | | |
| 18-34 | 56 | 34.8 | 110 | 65.2 | 8.4 |
| 35-54 | 120 | 45.7 | 124 | 54.3 | 7.1 |
| ≥55 | 120 | 55.3 | 96 | 44.7 | 7.2 |
| Education | | | | | |
| ≤High School | 108 | 56.7 | 77 | 43.3 | 8.1 |
| >High School | 194 | 40.4 | 262 | 59.6 | 5.1 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

# Current Smoker

## *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 120 | 16.8 | 522 | 83.2 | 3.2 |
| Gender | | | | | |
| Male | 47 | 16.9 | 206 | 83.1 | 5.0 |
| Female | 73 | 16.7 | 316 | 83.3 | 4.1 |
| Age | | | | | |
| 18-34 | 40 | 24.6 | 126 | 75.4 | 7.4 |
| 35-54 | 52 | 16.8 | 192 | 83.2 | 4.8 |
| ≥55 | 27 | 9.8 | 189 | 90.2 | 3.8 |
| Education | | | | | |
| ≤High School | 56 | 26.7 | 129 | 73.3 | 7.0 |
| >High School | 64 | 12.7 | 392 | 87.3 | 3.4 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Status of Cigarette Smoking

*Albany*

| Demographic Groups | n[1] | Everyday Smoker %[2] | C.I.[3] | n | Someday Smoker % | C.I. | n | Former or Never Smoked % | C.I. |
|---|---|---|---|---|---|---|---|---|---|
| Total | 91 | 12.8 | 2.9 | 29 | 4.0 | 1.6 | 522 | 83.2 | 3.2 |
| Gender | | | | | | | | | |
| Male | 37 | 13.6 | 4.6 | 10 | 3.4 | 2.4 | 206 | 83.1 | 5.0 |
| Female | 54 | 12.2 | 3.6 | 19 | 4.5 | 2.1 | 316 | 83.3 | 4.1 |
| Age | | | | | | | | | |
| 18-34 | 29 | 17.6 | 6.6 | 11 | 7.0 | 4.2 | 126 | 75.4 | 7.4 |
| 35-54 | 39 | 13.4 | 4.5 | 13 | 3.5 | 1.9 | 192 | 83.2 | 4.8 |
| ≥55 | 22 | 8.1 | 3.6 | 5 | 1.7 | 1.5 | 189 | 90.2 | 3.8 |
| Education | | | | | | | | | |
| ≤High School | 48 | 21.7 | 6.3 | 8 | 5.0 | 3.8 | 129 | 73.3 | 7.0 |
| >High School | 43 | 9.1 | 3.1 | 21 | 3.5 | 1.6 | 392 | 87.3 | 3.4 |

Expanded BRFSS, 2003 - New York State Department of Health

# Home Smoking Policies

## *Albany*

| Demographic Groups | n[1] | Not Allowed %[2] | C.I.[3] | n | Allowed Some Places % | C.I. | n | Allowed Anywhere % | C.I. |
|---|---|---|---|---|---|---|---|---|---|
| Total | 429 | 66.7 | 4.3 | 80 | 13.7 | 3.2 | 133 | 19.7 | 3.6 |
| Gender | | | | | | | | | |
| Male | 165 | 66.0 | 6.6 | 30 | 12.4 | 4.6 | 58 | 21.6 | 5.8 |
| Female | 264 | 67.3 | 5.5 | 50 | 14.7 | 4.4 | 75 | 18.0 | 4.4 |
| Age | | | | | | | | | |
| 18-34 | 111 | 61.8 | 8.8 | 29 | 21.7 | 7.6 | 26 | 16.5 | 7.2 |
| 35-54 | 167 | 70.8 | 6.5 | 30 | 10.4 | 3.9 | 47 | 18.8 | 5.8 |
| ≥55 | 142 | 67.4 | 6.7 | 20 | 9.6 | 4.3 | 54 | 23.0 | 5.9 |
| Education | | | | | | | | | |
| ≤High School | 101 | 54.7 | 8.2 | 29 | 16.9 | 6.5 | 55 | 28.5 | 7.3 |
| >High School | 327 | 71.7 | 4.9 | 51 | 12.3 | 3.6 | 78 | 16.0 | 4.1 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Home Smoking

*Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 143 | 23.9 | 500 | 76.1 | 4.0 |
| Gender | | | | | |
| Male | 63 | 25.5 | 190 | 74.5 | 6.3 |
| Female | 80 | 22.5 | 310 | 77.5 | 5.1 |
| Age | | | | | |
| 18-34 | 48 | 35.1 | 118 | 64.9 | 8.8 |
| 35-54 | 55 | 21.6 | 189 | 78.4 | 6.0 |
| ≥55 | 39 | 16.5 | 178 | 83.5 | 5.1 |
| Education | | | | | |
| ≤High School | 58 | 32.9 | 128 | 67.1 | 7.9 |
| >High School | 85 | 20.1 | 371 | 79.9 | 4.6 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)
### *Albany*

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---|---|---|---|---|---|
| Total | 80 | 14.5 | 553 | 85.5 | 3.5 |
| Gender | | | | | |
| Male | 51 | 21.8 | 198 | 78.2 | 6.2 |
| Female | 29 | 8.1 | 355 | 91.9 | 3.5 |
| Age | | | | | |
| 18-34 | 42 | 28.9 | 123 | 71.1 | 8.6 |
| 35-54 | 29 | 11.3 | 211 | 88.7 | 4.3 |
| ≥55 | 9 | 4.3 | 204 | 95.7 | 3.0 |
| Education | | | | | |
| ≤High School | 19 | 11.3 | 163 | 88.7 | 5.2 |
| >High School | 61 | 15.8 | 389 | 84.2 | 4.5 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution
²Weighted Percent
³95% Confidence Interval

## At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)

### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 40 | 8.3 | 590 | 91.7 | 3.0 |
| Gender | | | | | |
| Male | 17 | 9.8 | 229 | 90.2 | 5.2 |
| Female | 23 | 7.0 | 361 | 93.0 | 3.4 |
| Age | | | | | |
| 18-34 | 20 | 17.3 | 144 | 82.7 | 8.0 |
| 35-54 | 8 | 2.4 | 230 | 97.6 | 1.7 |
| ≥55 | 12 | 6.0 | 201 | 94.0 | 3.5 |
| Education | | | | | |
| ≤High School | 14 | 7.3 | 167 | 92.7 | 4.2 |
| >High School | 26 | 8.7 | 422 | 91.3 | 3.9 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Overweight or Obese, BMI(Body Mass Index) 25 or More
### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 330 | 54.0 | 285 | 46.0 | 4.5 |
| Gender | | | | | |
| Male | 167 | 65.2 | 84 | 34.8 | 6.6 |
| Female | 163 | 43.4 | 201 | 56.6 | 5.8 |
| Age | | | | | |
| 18-34 | 67 | 38.2 | 92 | 61.8 | 8.2 |
| 35-54 | 134 | 62.0 | 102 | 38.0 | 6.8 |
| ≥55 | 126 | 61.2 | 84 | 38.8 | 7.1 |
| Education | | | | | |
| ≤High School | 96 | 53.8 | 82 | 46.2 | 8.3 |
| >High School | 234 | 54.1 | 202 | 45.9 | 5.3 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+)
### *Albany*

| Demographic Groups | n[1] | Underweight\Normal %[2] | C.I.[3] | n | Overweight % | C.I. | n | Obese % | C.I. |
|---|---|---|---|---|---|---|---|---|---|
| Total | 285 | 46.1 | 4.5 | 208 | 34.7 | 4.2 | 121 | 19.2 | 3.5 |
| Gender | | | | | | | | | |
| Male | 84 | 34.8 | 6.6 | 111 | 44.7 | 6.7 | 56 | 20.5 | 5.2 |
| Female | 201 | 56.7 | 5.8 | 97 | 25.3 | 4.9 | 65 | 18.0 | 4.6 |
| Age | | | | | | | | | |
| 18-34 | 92 | 61.8 | 8.2 | 42 | 24.8 | 7.1 | 25 | 13.4 | 5.2 |
| 35-54 | 102 | 38.0 | 6.8 | 84 | 40.4 | 7.2 | 50 | 21.7 | 6.2 |
| ≥55 | 84 | 38.9 | 7.1 | 80 | 38.3 | 7.1 | 45 | 22.8 | 6.4 |
| Education | | | | | | | | | |
| ≤High School | 82 | 46.3 | 8.4 | 55 | 30.0 | 7.4 | 40 | 23.7 | 7.3 |
| >High School | 202 | 45.9 | 5.3 | 153 | 36.7 | 5.1 | 81 | 17.4 | 3.8 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Ever Had Mammography Screening, Women Aged 40 Years and Older
### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Female | 236 | 95.2 | 12 | 4.8 | 2.8 |
| Age | | | | | |
| 40-64 | 165 | 95.7 | 8 | 4.3 | 3.1 |
| ≥65 | 71 | 94.4 | 4 | 5.6 | 5.6 |
| Education | | | | | |
| ≤High School | 81 | 94.5 | 6 | 5.5 | 4.5 |
| >High School | 155 | 95.6 | 6 | 4.4 | 3.6 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

# Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older

## *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Female | 207 | 84.9 | 39 | 15.1 | 4.7 |
| Age | | | | | |
| 40-64 | 148 | 87.4 | 24 | 12.6 | 5.1 |
| ≥65 | 59 | 80.2 | 15 | 19.8 | 9.5 |
| Education | | | | | |
| ≤High School | 68 | 82.8 | 18 | 17.2 | 7.9 |
| >High School | 139 | 86.1 | 21 | 13.9 | 5.9 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger**

*Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 89 | 19.9 | 412 | 80.1 | 4.5 |
| Gender | | | | | |
| Male | 20 | 12.8 | 187 | 87.2 | 6.3 |
| Female | 69 | 26.9 | 225 | 73.1 | 6.3 |
| Age | | | | | |
| 18-34 | 59 | 38.4 | 105 | 61.6 | 8.8 |
| 35-54 | 27 | 9.3 | 210 | 90.7 | 4.2 |
| 55-64 | 3 | 2.8 | 97 | 97.2 | 3.5 |
| Education | | | | | |
| ≤High School | 33 | 29.1 | 97 | 70.9 | 9.9 |
| >High School | 56 | 16.6 | 315 | 83.4 | 4.9 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Sexually Active, Age 18-64 Years Old

### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 369 | 85.8 | 91 | 14.2 | 3.4 |
| Gender | | | | | |
| Male | 165 | 88.6 | 30 | 11.4 | 4.7 |
| Female | 204 | 82.9 | 61 | 17.1 | 4.9 |
| Age | | | | | |
| 18-34 | 144 | 91.6 | 15 | 8.4 | 5.0 |
| 35-54 | 167 | 84.7 | 46 | 15.3 | 5.1 |
| 55-64 | 58 | 73.9 | 30 | 26.1 | 9.4 |
| Education | | | | | |
| ≤High School | 86 | 74.0 | 34 | 26.0 | 9.4 |
| >High School | 283 | 90.0 | 57 | 10.0 | 2.8 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger
### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 87 | 21.5 | 296 | 78.5 | 4.7 |
| Gender | | | | | |
| Male | 42 | 21.0 | 129 | 79.0 | 6.6 |
| Female | 45 | 22.0 | 167 | 78.0 | 6.8 |
| Age | | | | | |
| 18-34 | 47 | 31.2 | 102 | 68.8 | 8.6 |
| 35-54 | 35 | 16.8 | 136 | 83.2 | 5.8 |
| 55-64 | 5 | 5.3 | 58 | 94.7 | 4.6 |
| Education | | | | | |
| ≤High School | 18 | 19.9 | 71 | 80.1 | 9.8 |
| >High School | 69 | 22.0 | 225 | 78.0 | 5.4 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

# Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)
## *Albany*

| Demographic Groups | n[1] | Use BC %[2] | C.I.[3] | n | No BC at risk % | C.I. | n | No BC % | C.I. |
|---|---|---|---|---|---|---|---|---|---|
| Total | 207 | 62.5 | 6.0 | 47 | 15.1 | 4.6 | 82 | 22.5 | 5.1 |
| Gender | | | | | | | | | |
| Male | 103 | 60.1 | 8.3 | 25 | 16.5 | 6.8 | 47 | 23.4 | 7.0 |
| Female | 104 | 65.8 | 8.4 | 22 | 13.0 | 5.5 | 35 | 21.2 | 7.5 |
| Age | | | | | | | | | |
| 18-34 | 107 | 63.0 | 8.8 | 24 | 16.7 | 7.0 | 32 | 20.3 | 7.6 |
| 35-59 | 100 | 61.9 | 7.8 | 23 | 13.0 | 5.5 | 50 | 25.1 | 6.7 |
| Education | | | | | | | | | |
| ≤High School | 39 | 52.3 | 12.5 | 13 | 16.5 | 8.9 | 27 | 31.2 | 11.7 |
| >High School | 168 | 65.6 | 6.8 | 34 | 14.6 | 5.4 | 55 | 19.8 | 5.6 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Physician Diagnosed Heart Attack, Angina or Stroke
### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 51 | 7.9 | 583 | 92.1 | 2.3 |
| Gender | | | | | |
| Male | 32 | 10.8 | 220 | 89.2 | 3.8 |
| Female | 19 | 5.4 | 363 | 94.6 | 2.7 |
| Age | | | | | |
| 18-64 | 25 | 4.9 | 483 | 95.1 | 2.1 |
| ≥65 | 26 | 22.2 | 89 | 77.8 | 8.1 |
| Education | | | | | |
| ≤High School | 23 | 13.7 | 161 | 86.3 | 5.7 |
| >High School | 27 | 5.3 | 422 | 94.7 | 2.1 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina)
### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 39 | 6.2 | 594 | 93.8 | 2.0 |
| Gender | | | | | |
| Male | 26 | 9.2 | 225 | 90.8 | 3.6 |
| Female | 13 | 3.6 | 369 | 96.4 | 2.1 |
| Age | | | | | |
| 18-64 | 17 | 3.2 | 490 | 96.8 | 1.6 |
| ≥65 | 22 | 20.0 | 93 | 80.0 | 8.0 |
| Education | | | | | |
| ≤High School | 20 | 11.3 | 163 | 88.7 | 5.1 |
| >High School | 18 | 3.9 | 431 | 96.1 | 1.9 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older
### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Male | 96 | 64.4 | 58 | 35.6 | 8.0 |
| Age | | | | | |
| 40-64 | 64 | 56.7 | 51 | 43.3 | 9.7 |
| ≥65 | 32 | 86.3 | 7 | 13.7 | 10.3 |
| Education | | | | | |
| ≤High School | 28 | 63.7 | 16 | 36.3 | 15.1 |
| >High School | 68 | 65.0 | 41 | 35.0 | 9.4 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

### Albany

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Male | 86 | 59.1 | 66 | 40.9 | 8.3 |
| Age | | | | | |
| 40-64 | 55 | 50.1 | 58 | 49.9 | 10.0 |
| ≥65 | 31 | 84.7 | 8 | 15.3 | 10.7 |
| Education | | | | | |
| ≤High School | 26 | 59.4 | 18 | 40.6 | 15.4 |
| >High School | 60 | 59.3 | 47 | 40.7 | 9.8 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Ever Had a Digital Rectal Exam, Men Aged 40 and Older
### Albany

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Male | 137 | 87.1 | 19 | 12.9 | 5.7 |
| Age | | | | | |
| 40-64 | 100 | 85.3 | 16 | 14.7 | 7.1 |
| ≥65 | 37 | 92.3 | 3 | 7.7 | 8.6 |
| Education | | | | | |
| ≤High School | 38 | 83.1 | 8 | 16.9 | 11.3 |
| >High School | 98 | 88.7 | 11 | 11.3 | 6.6 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Ever Used a Blood Stool Test at Home, Aged 50 and Older
### *Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 120 | 42.6 | 152 | 57.4 | 6.5 |
| Gender | | | | | |
| Male | 44 | 41.3 | 62 | 58.7 | 10.0 |
| Female | 76 | 43.6 | 90 | 56.4 | 8.4 |
| Age | | | | | |
| 50-64 | 72 | 44.0 | 87 | 56.0 | 8.7 |
| ≥65 | 48 | 40.7 | 65 | 59.3 | 9.6 |
| Education | | | | | |
| ≤High School | 41 | 41.0 | 62 | 59.0 | 10.5 |
| >High School | 79 | 43.8 | 89 | 56.2 | 8.2 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older

*Albany*

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 50 | 18.8 | 218 | 81.2 | 5.1 |
| Gender | | | | | |
| Male | 18 | 15.8 | 88 | 84.2 | 7.3 |
| Female | 32 | 21.3 | 130 | 78.7 | 7.2 |
| Age | | | | | |
| 50-64 | 30 | 20.0 | 128 | 80.0 | 7.1 |
| ≥65 | 20 | 17.2 | 90 | 82.8 | 7.4 |
| Education | | | | | |
| ≤High School | 26 | 29.0 | 76 | 71.0 | 10.0 |
| >High School | 24 | 12.2 | 141 | 87.8 | 5.0 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older
### Albany

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 73 | 27.3 | 195 | 72.7 | 5.9 |
| Gender | | | | | |
| Male | 29 | 26.3 | 77 | 73.7 | 8.8 |
| Female | 44 | 28.3 | 118 | 71.7 | 7.8 |
| Age | | | | | |
| 50-64 | 42 | 28.1 | 116 | 71.9 | 7.9 |
| ≥65 | 31 | 26.4 | 79 | 73.6 | 8.7 |
| Education | | | | | |
| ≤High School | 32 | 33.2 | 70 | 66.8 | 10.2 |
| >High School | 41 | 23.6 | 124 | 76.4 | 7.0 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older
### Albany

| Demographic Groups | n[1] | Yes %[2] | n | No % | C.I.[3] |
|---|---|---|---|---|---|
| Total | 167 | 63.2 | 105 | 36.8 | 6.2 |
| Gender | | | | | |
| Male | 74 | 72.3 | 31 | 27.7 | 9.0 |
| Female | 93 | 55.6 | 74 | 44.4 | 8.4 |
| Age | | | | | |
| 50-64 | 96 | 61.4 | 61 | 38.6 | 8.6 |
| ≥65 | 71 | 65.3 | 44 | 34.7 | 9.1 |
| Education | | | | | |
| ≤High School | 55 | 59.0 | 48 | 41.0 | 10.3 |
| >High School | 112 | 66.2 | 56 | 33.8 | 7.9 |

[1]Percents based on row denominators of less than 50 are unstable and should be used with caution
[2]Weighted Percent
[3]95% Confidence Interval

## Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older
### *Albany*

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---|---|---|---|---|---|
| Total | 158 | 59.8 | 114 | 40.2 | 6.4 |
| Gender | | | | | |
| Male | 71 | 69.9 | 34 | 30.1 | 9.2 |
| Female | 87 | 51.4 | 80 | 48.6 | 8.5 |
| Age | | | | | |
| 50-64 | 92 | 58.7 | 65 | 41.3 | 8.7 |
| ≥65 | 66 | 61.1 | 49 | 38.9 | 9.3 |
| Education | | | | | |
| ≤High School | 52 | 55.8 | 51 | 44.2 | 10.5 |
| >High School | 106 | 62.7 | 62 | 37.3 | 8.0 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution
²Weighted Percent
³95% Confidence Interval

**Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older**

*Albany*

| Demographic Groups | n¹ | Yes %² | n | No % | C.I.³ |
|---|---|---|---|---|---|
| Total | 175 | 66.9 | 99 | 33.1 | 6.0 |
| Gender | | | | | |
| Male | 76 | 74.7 | 30 | 25.3 | 8.5 |
| Female | 99 | 60.4 | 69 | 39.6 | 8.2 |
| Age | | | | | |
| 50-64 | 101 | 66.5 | 58 | 33.5 | 8.0 |
| ≥65 | 74 | 67.5 | 41 | 32.5 | 8.9 |
| Education | | | | | |
| ≤High School | 63 | 68.1 | 41 | 31.9 | 9.3 |
| >High School | 112 | 66.5 | 57 | 33.5 | 7.8 |

¹Percents based on row denominators of less than 50 are unstable and should be used with caution
²Weighted Percent
³95% Confidence Interval