

Expanded BRFSS, 2003

Queens Locality

Table of Contents:

Optional Module 1: Adult Asthma History 1
Optional Module 5: Diabetes..... 2
Optional Module 7: Disability/Quality of Life..... 6

Optional Module 10: Health Care Coverage 10
Optional Module 13: Injury Control - Falls..... 12
Optional Module 19: Tobacco Consumption 13
Optional Module 20: Tobacco - Media 14
Optional Module 21: Tobacco – other products 22
Optional Module 22: Tobacco – Work site ETS 23
Optional Module 23: Seat Belts..... 25
Optional Module 24: Weight Control..... 27

New York State Department of Health

Expanded BRFSS, 2003 - New York State Department of Health

Have had an Asthma Episode or Attack in the Past 12 Months, among Those with Asthma

Queens

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	20	41.3	23	58.7	17.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Check Blood for Glucose or Sugar at Least Once per Day among Those with Diabetes

Queens

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	22	39.9	26	60.1	15.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Check Feet for Sores or Irritations at Least Once per Day among Those with Diabetes

Queens

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	31	57.6	18	42.4	16.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had Feet Checked by a Medical Professional in Past Twelve Months, among Those with Diabetes

Queens

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	31	56.2	16	43.8	16.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had Dilated Eye Exam in Past 12 Months, among Those with Diabetes

Queens

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	32	73.5	14	26.5	14.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Adults with a Disability

Queens

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	96	13.9	487	86.1	3.0
Gender					
Male	38	13.2	197	86.8	4.5
Female	58	14.5	290	85.5	4.0
Age					
18-34	15	6.9	158	93.1	3.9
35-54	25	9.5	187	90.5	4.1
≥55	54	28.0	132	72.0	7.3
Education					
≤High School	40	15.9	174	84.1	5.4
>High School	56	12.9	304	87.1	3.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Adults with a Disability Requiring Help

Queens

Demographic Groups	n ¹	Disability need help % ²	C.I. ³	n	Disability no help needed %	C.I.	n	No Disability %	C.I.
Total	38	5.6	1.9	58	8.3	2.4	487	86.1	3.0
Gender									
Male	10	4.0	2.7	28	9.1	3.7	197	86.8	4.5
Female	28	7.0	2.7	30	7.5	3.1	290	85.5	4.0
Age									
18-64	29	5.0	2.0	39	6.2	2.2	400	88.8	2.9
≥65	9	9.1	6.1	17	18.0	8.8	77	72.9	9.8
Education									
≤High School	17	5.8	3.2	23	10.1	4.5	174	84.1	5.4
>High School	21	5.6	2.5	35	7.2	2.6	304	87.1	3.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

This page left intentionally blank

This page left intentionally blank

Expanded BRFSS, 2003 - New York State Department of Health

Main Reason for Not Having Health Care Coverage

Queens

Demographic Groups	n ¹	Changed or lost job % ²	C.I. ³	n	Could not afford %	C.I.	n	Other %	C.I.
Total	10	13.6	9.7	20	20.8	9.6	60	65.7	11.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

No Health Care Coverage for Past Two Years Among Uninsured

Queens

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	37	40.0	55	60.0	11.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had a Fall in Past Three Months

Queens

Demographic Groups	n¹	No fall %²	C.I.³	n	Fall with injury %	C.I.	n	Fall no injury %	C.I.
Total	227	85.6	4.9	8	3.1	2.5	31	11.4	4.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Number of Cigarettes Smoked per day among Everyday Smokers

Queens

Demographic Groups	n¹	1-10 cigarettes %²	n	≥11 cigarettes %	C.I.³
Total	29	64.0	19	36.0	15.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Television

Queens

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	349	63.6	214	36.4	4.7
Gender					
Male	143	64.6	86	35.4	7.4
Female	206	62.8	128	37.2	5.9
Age					
18-34	125	72.5	46	27.5	7.9
35-54	129	63.4	79	36.6	7.7
≥55	88	52.7	85	47.3	8.6
Education					
≤High School	129	63.6	78	36.4	7.9
>High School	214	63.1	133	36.9	5.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Radio

Queens

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	184	33.5	375	66.5	4.6
Gender					
Male	87	38.2	142	61.8	7.4
Female	97	29.4	233	70.6	5.6
Age					
18-34	67	40.2	105	59.8	8.7
35-54	65	32.8	140	67.2	7.4
≥55	50	27.5	121	72.5	7.4
Education					
≤High School	80	37.1	126	62.9	7.6
>High School	102	31.0	242	69.0	5.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Billboards

Queens

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	189	37.7	365	62.3	4.9
Gender					
Male	76	39.1	153	60.9	7.6
Female	113	36.3	212	63.7	6.1
Age					
18-34	75	47.0	92	53.0	9.0
35-54	81	44.8	122	55.2	8.2
≥55	29	17.9	144	82.1	6.8
Education					
≤High School	69	38.2	134	61.8	8.1
>High School	117	37.2	226	62.8	6.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Buses

Queens

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	157	30.0	401	70.0	4.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Magazines

Queens

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	290	53.7	273	46.3	4.9
Gender					
Male	118	54.0	112	46.0	7.6
Female	172	53.4	161	46.6	6.2
Age					
18-34	87	50.8	84	49.2	8.9
35-54	116	59.4	90	40.6	7.8
≥55	81	49.0	95	51.0	8.5
Education					
≤High School	109	57.1	95	42.9	8.0
>High School	178	51.7	172	48.3	6.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from a Website

Queens

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	70	13.6	495	86.4	3.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Movie Theater

Queens

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	69	13.7	496	86.3	3.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard or Seen Anti-Tobacco Advertisement that Refers to the NYS Quitline in Past 30 Days

Queens

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	240	47.7	302	52.3	5.0
Gender					
Male	103	50.9	122	49.1	7.7
Female	137	44.7	180	55.3	6.4
Age					
18-34	78	50.9	92	49.1	8.9
35-54	92	49.3	104	50.7	8.2
≥55	64	40.0	102	60.0	8.5
Education					
≤High School	93	52.8	108	47.2	8.1
>High School	143	44.0	189	56.0	6.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Used or Tried Any Smokeless Tobacco Products such as Chewing Tobacco or Snuff

Queens

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	20	3.1	550	96.9	1.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Work Indoors Most of the Time

Queens

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	264	74.5	71	25.5	6.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Smoking Policy at Place of Work among Indoor Workers*Queens*

Demographic Groups	n¹	Not allowed anywhere %²	C.I.³	n	Allowed in some or all places %	C.I.	n	No official policy %	C.I.
Total	235	88.6	4.9	13	6.6	4.2	12	4.8	2.8
Gender									
Male	85	87.4	7.2	6	6.7	5.8	7	5.9	4.6
Female	150	89.7	6.6	7	6.4	5.9	5	3.9	3.5
Age									
18-34	77	84.4	10.4	5	11.3	9.8	4	4.3	4.5
35-54	107	88.3	6.2	6	4.7	4.0	8	7.0	5.0
≥55	47	99.0	1.9	1	1.0	1.9	0	0.0	0.0
Education									
≤High School	57	85.5	11.2	6	12.1	10.8	2	2.4	3.6
>High School	172	89.8	5.2	7	4.4	3.7	9	5.8	3.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution²Weighted Percent³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Use Seatbelt when Ride or Drive in a Car (Always, Nearly Always or Never Ride in a Car)

Queens

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	514	92.0	50	8.0	2.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Use Seatbelt when Ride or Drive in a Car (Always or Never ride in a Car)

Queens

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	443	78.1	121	21.9	4.1
Gender					
Male	173	73.8	57	26.2	6.9
Female	270	82.0	64	18.0	4.5
Age					
18-64	357	77.8	99	22.2	4.6
≥65	77	80.6	20	19.4	8.3
Education					
≤High School	155	75.8	52	24.2	6.8
>High School	281	79.6	67	20.4	5.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Trying to Lose Weight

Queens

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	283	48.4	281	51.6	4.9
Gender					
Male	102	42.9	127	57.1	7.6
Female	181	53.3	154	46.7	6.2
Age					
18-34	80	44.1	91	55.9	8.8
35-54	124	58.7	82	41.3	7.9
≥55	76	42.3	100	57.7	8.4
Education					
≤High School	105	50.2	101	49.8	8.1
>High School	173	46.5	176	53.5	6.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Trying to Maintain Current Weight of Those not Trying to Lose Weight

Queens

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	163	58.5	119	41.5	6.7
Gender					
Male	67	53.6	60	46.4	10.1
Female	96	63.9	59	36.1	8.4
Age					
18-34	45	45.8	46	54.2	11.8
35-54	50	68.9	32	31.1	10.5
≥55	64	65.5	37	34.5	10.3
Education					
≤High School	59	60.0	43	40.0	10.9
>High School	103	59.1	73	40.9	8.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Eating Fewer Calories or Less Fat to Lose or Maintain Weight

Queens

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	351	79.6	84	20.4	4.9
Gender					
Male	127	75.5	38	24.5	8.4
Female	224	82.7	46	17.3	5.5
Age					
18-34	93	75.1	31	24.9	9.4
35-54	143	81.4	29	18.6	8.1
≥55	110	81.9	23	18.1	7.6
Education					
≤High School	135	82.9	24	17.1	7.7
>High School	213	78.7	57	21.3	5.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Using Physical Exercise to Lose or Maintain Weight

Queens

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	290	65.9	156	34.1	5.2
Gender					
Male	102	63.1	67	36.9	8.7
Female	188	68.2	89	31.8	6.2
Age					
18-34	94	78.9	31	21.1	7.7
35-54	121	67.3	53	32.7	8.8
≥55	69	48.9	71	51.1	9.6
Education					
≤High School	91	58.7	73	41.3	8.9
>High School	194	70.8	82	29.2	6.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval